DAMA/LIBRA
results and perspectives

V. Caracciolo
National Laboratory of Gran Sasso

BLED-16
July 11-19, 2016

DAMA set-ups

an observatory for rare processes @ LNGS



- DAMA/LIBRA (DAMA/NaI)
- DAMA/LXe
- DAMA/R&D
- DAMA/Crys
- DAMA/Ge

Collaboration:

Roma Tor Vergata, Roma La Sapienza, LNGS, IHEP/Beijing
+ by-products and small scale expts.: INR-Kiev + other institutions
+ neutron meas.: ENEA-Frascati
+ in some studies on ββ decays (DST-MAE and Inter-Universities project):
IIT Kharagpur and Ropar, India

web site: http://people.roma2.infn.it/dama

Some direct detection processes:

- Scatterings on nuclei

  → detection of nuclear recoil energy



- Inelastic Dark Matter: **W + N → W* + N**

  → W has 2 mass states χ^+ , χ^- with δ mass splitting

  → Kinematical constraint for the inelastic scattering of χ^- on a nucleus

$$\frac{1}{2}\mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

- Excitation of bound electrons in scatterings on nuclei

  → detection of recoil nuclei + e.m. radiation

- Conversion of particle into e.m. radiation

  → detection of γ, X-rays, e^-



- Interaction only on atomic electrons

  → detection of e.m. radiation



... even WIMPs

- Interaction of light DMp (LDM) on e^- or nucleus with production of a lighter particle

  → detection of electron/nucleus recoil energy

e.g. sterile ν



e.g. signals from these candidates are **completely lost** in experiments based on "rejection procedures" of the e.m. component of their rate

… also other ideas …

The annual modulation: a model independent signature for the investigation of DM particles component in the galactic halo

With the present technology, the annual modulation is the main model independent signature for the DM signal. Although the modulation effect is expected to be relatively small a suitable large-mass, low-radioactive set-up with an efficient control of the running conditions can point out its presence.

Requirements:

1) Modulated rate according cosine
2) In low energy range
3) With a proper period (1 year)
4) With proper phase (about 2 June)
5) Just for single hit events in a multi-detector set-up
6) With modulation amplitude in the region of maximal sensitivity must be <7% for usually adopted halo distributions, but it can be larger in case of some possible scenarios

Drukier, Freese, Spergel PRD86; Freese et al. PRD88



- v_{sun} ~ 232 km/s (Sun vel in the halo)
- v_{orb} = 30 km/s (Earth vel around the Sun)
- $\gamma = \pi/3$, $\omega = 2\pi/T$, T = 1 year
- t_0 = 2nd June (when $v_\oplus$ is maximum)

$$v_\oplus(t) = v_{sun} + v_{orb}\cos\gamma\cos[\omega(t-t_0)]$$

$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k}\cos[\omega(t-t_0)]$$

the DM annual modulation signature has a different origin and peculiarities (e.g. the phase) than those effects correlated with the seasons

To mimic this signature, spurious effects and side reactions must not only - obviously - be able to account for the whole observed modulation amplitude, but also to satisfy contemporaneously all the requirements

The pioneer DAMA/NaI: ≈100 kg highly radiopure NaI(Tl)

Performances:

N.Cim.A112(1999)545-575, EPJC18(2000)283,
Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127

Results on rare processes:

- Possible Pauli exclusion principle violation — PLB408(1997)439
- CNC processes — PRC60(1999)065501
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell) — PLB460(1999)235
- Search for solar axions — PLB515(2001)6
- Exotic Matter search — EPJdirect C14(2002)1
- Search for superdense nuclear matter — EPJA23(2005)7
- Search for heavy clusters decays — EPJA24(2005)51

Results on DM particles:

- PSD — PLB389(1996)757
- Investigation on diurnal effect — N.Cim.A112(1999)1541
- Exotic Dark Matter search — PRL83(1999)4918
- **Annual Modulation Signature** — PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512, PLB480(2000)23, EPJC18(2000)283, PLB509(2001)197, EPJC23(2002)61, PRD66(2002)043503, Riv.N.Cim.26 n.1 (2003)1, IJMPD13(2004)2127, IJMPA21(2006)1445, EPJC47(2006)263, IJMPA22(2007)3155, EPJC53(2008)205, PRD77(2008)023506, MPLA23(2008)2125



Model independent evidence of a particle DM component in the galactic halo at 6.3σ C.L.

total exposure (7 annual cycles) 0.29 ton × yr

The DAMA/LIBRA set-up ~250 kg NaI(Tl)
(Large sodium Iodide Bulk for RAre processes)

As a result of a 2nd generation R&D for more radiopure NaI(Tl) by exploiting new chemical/physical radiopurification techniques
(all operations involving - including photos - in HP Nitrogen atmosphere)



Residual contaminations in the new DAMA/LIBRA NaI(Tl) detectors: ^{232}Th, ^{238}U and ^{40}K at level of 10^{-12} g/g

- Radiopurity, performances, procedures, etc.: NIMA592(2008)297, JINST 7 (2012) 03009
- Results on DM particles, Annual Modulation Signature: EPJC56(2008)333, EPJC67(2010)39, EPJC73(2013)2648.
  Related results: PRD84(2011)055014, EPJC72(2012)2064, IJMPA28(2013)1330022, EPJC74(2014)2827, EPJC74(2014)3196, EPJC75(2015)239, EPJC75(2015)400
- Results on rare processes: PEPv: EPJC62(2009)327; CNC: EPJC72(2012)1920; IPP in ^{241}Am: EPJA49(2013)64

The DAMA/LIBRA set-up

For details, radiopurity, performances, procedures, etc.
NIMA592(2008)297

Polyethylene/paraffin

- 25 x 9.7 kg NaI(Tl) in a 5x5 matrix
- two Suprasil-B light guides directly coupled to each bare crystal
- two PMTs working in coincidence at the single ph. el. threshold



5.5-7.5 phe/keV



~ 1m concrete from GS rock

- Dismounting/Installing protocol (with "Scuba" system)
- All the materials selected for low radioactivity
- Multicomponent passive shield (>10 cm of Cu, 15 cm of Pb + Cd foils, 10/40 cm Polyethylene/paraffin, about 1 m concrete, mostly outside the installation)
- Three-level system to exclude Radon from the detectors
- Calibrations in the same running conditions as production runs
- Installation in air conditioning + huge heat capacity of shield
- Monitoring/alarm system; many parameters acquired with the production data
- Pulse shape recorded by Waveform Analyzer Acqiris DC270 (2chs per detector), 1 *Gsample/s*, 8 bit, bandwidth 250 MHz
- Data collected from low energy up to MeV region, despite the hardware optimization was done for the low energy

DAMA/LIBRA calibrations

Low energy: various external gamma sources (^{241}Am, ^{133}Ba) and internal X-rays or gamma's (^{40}K, ^{125}I, ^{129}I), routine calibrations with ^{241}Am



$$\frac{\sigma_{LE}}{E} = \frac{(0.448 \pm 0.035)}{\sqrt{E(keV)}} + (9.1 \pm 5.1) \cdot 10^{-3}$$



The curves superimposed to the experimental data have been obtained by simulations

High energy: external sources of gamma rays (e.g. ^{137}Cs, ^{60}Co and ^{133}Ba) and gamma rays of 1461 keV due to ^{40}K decays in an adjacent detector, tagged by the 3.2 keV X-rays



The signals (unlike low energy events) for high energy events are taken only from one PMT



$$\frac{\sigma_{HE}}{E} = \frac{(1.12 \pm 0.06)}{\sqrt{E(keV)}} + (17 \pm 23) \cdot 10^{-4}$$

Thus, here and hereafter keV means keV electron equivalent

Complete DAMA/LIBRA-phase1

	Period	Mass (kg)	Exposure (kg×day)	$(\alpha - \beta^2)$
DAMA/LIBRA-1	Sept. 9, 2003 - July 21, 2004	232.8	51405	0.562
DAMA/LIBRA-2	July 21, 2004 - Oct. 28, 2005	232.8	52597	0.467
DAMA/LIBRA-3	Oct. 28, 2005 - July 18, 2006	232.8	39445	0.591
DAMA/LIBRA-4	July 19, 2006 - July 17, 2007	232.8	49377	0.541
DAMA/LIBRA-5	July 17, 2007 - Aug. 29, 2008	232.8	66105	0.468
DAMA/LIBRA-6	Nov. 12, 2008 - Sept. 1, 2009	242.5	58768	0.519
DAMA/LIBRA-7	Sep. 1, 2009 - Sept. 8, 2010	242.5	62098	0.515
DAMA/LIBRA-phase1	Sept. 9, 2003 - Sept. 8, 2010		379795 = 1.04 ton×yr	0.518
DAMA/NaI + DAMA/LIBRA-phase1:			1.33 ton×yr	

a ton × yr experiment? done

- EPJC56(2008)333
- EPJC67(2010)39
- EPJC73(2013)2648

- calibrations: ≈96 Mevents from sources
- acceptance window eff: 95 Mevents (≈3.5 Mevents/keV)

DAMA/LIBRA-phase1:

- First upgrade on Sept 2008: replacement of some PMTs in HP N_2 atmosphere, new Digitizers (U1063A Acqiris 1GS/s 8-bit High-speed cPCI), new DAQ system with optical read-out installed

DAMA/LIBRA-phase2 (running):

- Second upgrade at end 2010: replacement of all the PMTs with higher Q.E. ones from dedicated developments
- commissioning on 2011

Goal: lowering the software energy threshold

- Fall 2012: new preamplifiers installed + special trigger modules. Other new components in the electronic chain in development

Model Independent Annual Modulation Result

DAMA/NaI + DAMA/LIBRA-phase1 Total exposure: 487526 kg×day = **1.33 ton×yr**

Single-hit residuals rate of scintillation events *vs* time in 2-6 keV

EPJC 56(2008)333, EPJC 67(2010)39, EPJC 73(2013)2648



continuous line: t_0 = 152.5 d, T =1.0 y

A=(0.0110±0.0012) cpd/kg/keV

χ^2/dof = 70.4/86 9.2 σ C.L.

Absence of modulation? No

χ^2/dof=154/87 P(A=0) = 1.3×10^{-5}

Fit with all the parameters free:

A = (0.0112 ± 0.0012) cpd/kg/keV

t_0 = (144±7) d - T = (0.998±0.002) y



No systematics or side reaction able to account for the measured modulation amplitude and to satisfy all the peculiarities of the signature

Comparison between **single hit residual rate (red points)** and **multiple hit residual rate (green points)**; Clear modulation in the single hit events; No modulation in the residual rate of the multiple hit events

A=-(0.0005±0.0004) cpd/kg/keV



This result offers an additional strong support for the presence of DM particles in the galactic halo further excluding any side effect either from hardware or from software procedures or from background

The data favor the presence of a modulated behaviour with all the proper features for DM particles in the galactic halo at about 9.2σ C.L.

DAMA/NaI & DAMA/LIBRA experiments main upgrades and improvements



PHASE2

Minimal upgrade in Fall

July 2000 new DAQ and new electronic chain installed (MULTIPLEXER removed, now one TD channel for each detector):
(i) TD VXI Tektronix;
(ii) Digital Unix DAQ system;
(iii) GPIB-CAMAC.

July 2002 DAMA/NaI data taking completed

On 2003 DAMA/LIBRA has begun first operations

Sept.-Oct. 2008 – DAMA/LIBRA upgrade:
① one detector recovered by replacing a broken PMT
② a new optimization of some PMTs and HVs performed
③ all the TD replaced with new ones (U1063A Acqiris 8-bit 1GS/s DC270 High-Speed cPCI Digitizers)
④ a new DAQ with optical read-out installed.

The second DAMA/LIBRA upgrade in Fall 2010:
Replacement of all the PMTs with higher Q.E. ones from dedicated developments
(+new preamp in Fall 2012 and other developments in progress)

DAMA/LIBRA-phase2 *in data taking*

Rate behaviour above 6 keV

- No Modulation above 6 keV



Mod. Ampl. (6-10 keV): cpd/kg/keV

(0.0016 ± 0.0031) DAMA/LIBRA-1
-(0.0010 ± 0.0034) DAMA/LIBRA-2
-(0.0001 ± 0.0031) DAMA/LIBRA-3
-(0.0006 ± 0.0029) DAMA/LIBRA-4
-(0.0021 ± 0.0026) DAMA/LIBRA-5
(0.0029 ± 0.0025) DAMA/LIBRA-6
-(0.0023 ± 0.0024) DAMA/LIBRA-7

→ statistically consistent with zero

DAMA/LIBRA-phase1



- No modulation in the whole energy spectrum:
  studying integral rate at higher energy, R_{90}
  - R_{90} percentage variations with respect to their mean values for single crystal in the DAMA/LIBRA running periods
  - Fitting the behaviour with time, adding a term modulated with period and phase as expected for DM particles:

Period	Mod. Ampl.
DAMA/LIBRA-1	-(0.05±0.19) cpd/kg
DAMA/LIBRA-2	-(0.12±0.19) cpd/kg
DAMA/LIBRA-3	-(0.13±0.18) cpd/kg
DAMA/LIBRA-4	(0.15±0.17) cpd/kg
DAMA/LIBRA-5	(0.20±0.18) cpd/kg
DAMA/LIBRA-6	-(0.20±0.16) cpd/kg
DAMA/LIBRA-7	-(0.28±0.18) cpd/kg

consistent with zero

$\sigma \approx 1\%$, fully accounted by statistical considerations

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region → R_{90} ~ tens cpd/kg → ~ 100 σ far away

No modulation above 6 keV
This accounts for all sources of bckg and is consistent with the studies on the various components

Energy distribution of the modulation amplitudes

The modulation amplitude, S_m, obtained by maximum likelihood method

$$R(t) = S_0 + S_m \cos\left[\omega\left(t - t_0\right)\right]$$

DAMA/NaI + DAMA/LIBRA-phase1

total exposure: 487526 kg×day ≈**1.33 ton×yr**



$T=2\pi/\omega$=1 yr

t_0= 152.5 day

A clear modulation is present in the (2-6) keV energy interval, while S_m values compatible with zero are present just above

The S_m values in the (6–20) keV energy interval have random fluctuations around zero with χ^2 equal to 35.8 for 28 degrees of freedom (upper tail probability 15%)

Is there a sinusoidal contribution in the signal? phase ≠ 152.5 day?

DAMA/NaI + DAMA/LIBRA-phase1

total exposure: 487526 kg×day ≈**1.33 ton×yr**

$$R(t) = S_0 + S_m \cos\left[\omega(t - t_0)\right] + Z_m \sin\left[\omega(t - t_0)\right] = S_0 + Y_m \cos\left[\omega(t - t^*)\right]$$

For Dark Matter signals:

- $|Z_m| \ll |S_m| \approx |Y_m|$
- $t^* \approx t_0 = 152.5d$
- $\omega = 2\pi/T$
- T = 1 year



Slight differences from 2nd June are expected in case of contributions from non thermalized DM components (as e.g. the SagDEG stream)



E (keV)	S_m (cpd/kg/keV)	Z_m (cpd/kg/keV)	Y_m (cpd/kg/keV)	t* (day)
2-6	0.0106 ± 0.0012	-0.0006 ± 0.0012	0.0107 ± 0.0012	149.5 ± 7.0
6-14	0.0001 ± 0.0007	0.0000 ± 0.0005	0.0001 ± 0.0008	--

No role for μ in DAMA annual modulation result

✓ **Direct μ interaction in DAMA/LIBRA set-up:**

DAMA/LIBRA surface ≈0.13 m^2

μ flux @ DAMA/LIBRA ≈2.5 μ/day

It cannot mimic the signature: already excluded by R_{90}, by *multi-hits* analysis + different phase, etc.



MonteCarlo simulation

✓ **Rate, R_n, of fast neutrons produced by μ:**

- Φ_μ @ LNGS ≈ 20 μ $m^{-2}d^{-1}$ (±1.5% modulated)
- Annual modulation amplitude at low energy due to μ **modulation**:

$$S_m^{(\mu)} = R_n\, g\, \varepsilon\, f_{\Delta E}\, f_{single}\, 2\% / (M_{setup}\, \Delta E)$$

$S_m^{(\mu)} < (0.3\text{-}2.4) \times 10^{-5}$ cpd/kg/keV

Moreover, this modulation also induces a variation in other parts of the energy spectrum and in the *multi-hits* events

It cannot mimic the signature: already excluded by R_{90}, by *multi-hits* analysis + different phase, etc.

✓ **Inconsistency of the phase between DAMA signal and μ modulation**

μ flux @ LNGS (MACRO, LVD, BOREXINO) ≈$3 \cdot 10^{-4}$ $m^{-2}s^{-1}$; modulation amplitude 1.5%; **phase**: July 7 ± 6 d, June 29 ± 6 d (Borexino)

The DAMA phase: May 26 ± 7 days (stable over 13 years)

The DAMA phase is 5.7σ far from the LVD/BOREXINO phases of muons (7.1 σ far from MACRO measured phase)



… many others arguments EPJC72(2012)2064, EPJC74(2014)3196

- Contributions to the total neutron flux at LNGS;
- Counting rate in DAMA/LIBRA for *single-hit* events, in the (2 – 6) keV energy region induced by:
  - neutrons,
  - muons,
  - solar neutrinos.

EPJC 74 (2014) 3196 (also EPJC 56 (2008) 333, EPJC 72 (2012) 2064,IJMPA 28 (2013) 1330022)

Modulation amplitudes

	Source	$\Phi_{0,k}^{(n)}$ (neutrons cm^{-2} s^{-1})	η_k	t_k	$R_{0,k}$ (cpd/kg/keV)		$A_k = R_{0,k}\eta_k$ (cpd/kg/keV)
SLOW neutrons	thermal n ($10^{-2} - 10^{-1}$ eV)	1.08×10^{-6} [15]	$\simeq 0$ however $\ll 0.1$ [2, 7, 8]	–	$< 8 \times 10^{-6}$	[2, 7, 8]	$\ll 8 \times 10^{-7}$
	epithermal n (eV-keV)	2×10^{-6} [15]	$\simeq 0$ however $\ll 0.1$ [2, 7, 8]	–	$< 3 \times 10^{-3}$	[2, 7, 8]	$\ll 3 \times 10^{-4}$
FAST neutrons	fission, $(\alpha, n) \rightarrow$ n (1-10 MeV)	$\simeq 0.9 \times 10^{-7}$ [17]	$\simeq 0$ however $\ll 0.1$ [2, 7, 8]	–	$< 6 \times 10^{-4}$	[2, 7, 8]	$\ll 6 \times 10^{-5}$
	$\mu \rightarrow$ n from rock (> 10 MeV)	$\simeq 3 \times 10^{-9}$ (see text and ref. [12])	0.0129 [23]	end of June [23, 7, 8]	$\ll 7 \times 10^{-4}$	(see text and [2, 7, 8])	$\ll 9 \times 10^{-6}$
	$\mu \rightarrow$ n from Pb shield (> 10 MeV)	$\simeq 6 \times 10^{-9}$ (see footnote 3)	0.0129 [23]	end of June [23, 7, 8]	$\ll 1.4 \times 10^{-3}$	(see text and footnote 3)	$\ll 2 \times 10^{-5}$
	$\nu \rightarrow$ n (few MeV)	$\simeq 3 \times 10^{-10}$ (see text)	0.03342 *	Jan. 4th *	$\ll 7 \times 10^{-5}$	(see text)	$\ll 2 \times 10^{-6}$
	direct μ	$\Phi_0^{(\mu)} \simeq 20\ \mu$ m^{-2}d^{-1} [20]	0.0129 [23]	end of June [23, 7, 8]	$\simeq 10^{-7}$	[2, 7, 8]	$\simeq 10^{-9}$
	direct ν	$\Phi_0^{(\nu)} \simeq 6 \times 10^{10}\ \nu$ cm^{-2}s^{-1} [26]	0.03342 *	Jan. 4th *	$\simeq 10^{-5}$	[31]	3×10^{-7}

* The annual modulation of solar neutrino is due to the different Sun-Earth distance along the year; so the relative modulation amplitude is twice the eccentricity of the Earth orbit and the phase is given by the perihelion.

All are negligible w.r.t. the annual modulation amplitude observed by DAMA/LIBRA and they cannot contribute to the observed modulation amplitude.

+ In no case neutrons (of whatever origin) can mimic the DM annual modulation signature since some of the **peculiar requirements of the signature** would fail, such as the neutrons would induce e.g. variations in all the energy spectrum, variation in the multiple hit events,... which were not observed.

Summary of the results obtained in the additional investigations of possible systematics or side reactions – DAMA/LIBRA-phase1

(NIMA592(2008)297, EPJC56(2008)333, J. Phys. Conf. ser. 203(2010)012040, arXiv:0912.0660, S.I.F.Atti Conf.103(211), Can. J. Phys. 89 (2011) 11, Phys.Proc.37(2012)1095, EPJC72(2012)2064, arxiv:1210.6199 & 1211.6346, IJMPA28(2013)1330022, EPJC74(2014)3196)

Source	Main comment	Cautious upper limit (90%C.L.)
RADON	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	**$<2.5\times10^{-6}$ cpd/kg/keV**
TEMPERATURE	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield→ huge heat capacity + T continuously recorded	**$<10^{-4}$ cpd/kg/keV**
NOISE	Effective full noise rejection near threshold	**$<10^{-4}$ cpd/kg/keV**
ENERGY SCALE	Routine + intrinsic calibrations	**$<1\text{-}2\times10^{-4}$ cpd/kg/keV**
EFFICIENCIES	Regularly measured by dedicated calibrations	**$<10^{-4}$ cpd/kg/keV**
BACKGROUND	No modulation above 6 keV; no modulation in the (2-6) keV *multiple-hits* events; this limit includes all possible sources of background	**$<10^{-4}$ cpd/kg/keV**
SIDE REACTIONS	Muon flux variation measured at LNGS	**$<3\times10^{-5}$ cpd/kg/keV**

+ they cannot satisfy all the requirements of annual modulation signature

Thus, they cannot mimic the observed annual modulation effect

Model-independent evidence by DAMA/NaI and DAMA/LIBRA

well compatible with several candidates in many astrophysical, nuclear and particle physics scenarios

Neutralino as LSP in various SUSY theories

Various kinds of WIMP candidates with several different kind of interactions Pure SI, pure SD, mixed + Migdal effect +channeling,... (from low to high mass)

a heavy ν of the 4-th family

Pseudoscalar, scalar or mixed light bosons with axion-like interactions

WIMP with preferred inelastic scattering

Mirror Dark Matter

Light Dark Matter

Self interacting Dark Matter

Dark Matter (including some scenarios for WIMP) electron-interacting

Sterile neutrino



heavy exotic canditates, as "4th family atoms", ...

Elementary Black holes such as the Daemons

Kaluza Klein particles

... and more

Is it an “universal” and “correct” way to approach the problem of DM and comparisons?



No, it isn’t. This is just a largely arbitrary/partial/incorrect exercise



About interpretation

See e.g.: Riv.N.Cim.26 n.1(2003)1, IJMPD13(2004)2127, EPJC47(2006)263, IJMPA21(2006)1445, EPJC56(2008)333, PRD84(2011)055014, IJMPA28(2013)1330022

…models…

- Which particle?
- Which interaction coupling?
- Which Form Factors for each target-material?
- Which Spin Factor?
- Which nuclear model framework?
- Which scaling law?
- Which halo model, profile and related parameters?
- Streams?
- …

…and experimental aspects…

- Exposures
- Energy threshold
- Detector response (phe/keV)
- Energy scale and energy resolution
- Calibrations
- Stability of all the operating conditions.
- Selections of detectors and of data.
- Subtraction/rejection procedures and stability in time of all the selected windows and related quantities
- Efficiencies
- Definition of fiducial volume and non-uniformity
- Quenching factors, channeling, …
- …

Uncertainty in experimental parameters, as well as necessary assumptions on various related astrophysical, nuclear and particle-physics aspects, affect all the results at various extent, both in terms of exclusion plots and in terms of allowed regions/volumes. Thus comparisons with a fixed set of assumptions and parameters' values are intrinsically strongly uncertain.

No experiment can be directly compared in model independent way with DAMA

... an example in literature...

Case of DM particles inducing elastic scatterings on target-nuclei, SI case



Regions in the nucleon cross section vs DM particle mass plane

- Some velocity distributions and uncertainties considered.
- The DAMA regions represent the domain where the likelihood-function values differ more than 7.5σ from the null hypothesis (absence of modulation).
- For CoGeNT a fixed value for the Ge quenching factor and a Helm form factor with fixed parameters are assumed.
- The CoGeNT region includes configurations whose likelihood-function values differ more than 1.64σ from the null hypothesis (absence of modulation). This corresponds roughly to 90% C.L. far from zero signal.

Including the Migdal effect
→Towards lower mass/higher σ

DAMA allowed regions for a particular set of astrophysical, nuclear and particle Physics assumptions without (green), with (blue) channeling, with energy-dependent Quenching Factors (red);

7.5 σ C.L.

PRD84(2011)055014, IJMPA28(2013)1330022



Co-rotating halo,
Non thermalized component
→ Enlarge allowed region towards larger mass

CoGeNT; qf at fixed assumed value

1.64 σ C.L.

Combining channeling and energy dependence of q.f. (AstrPhys33 (2010) 40)
→Towards lower σ

Other examples

Scratching Below the Surface of the Most General Parameter Space
(S. Scopel arXiv:1505.01926)

Most general approach: consider ALL possible NR couplings, including those depending on velocity and momentum

- A much wider parameter space opens up
- First explorations show that indeed large rooms for compatibility can be achieved

$$\mathcal{O}_1 = 1_\chi 1_N,$$
$$\mathcal{O}_2 = (v^\perp)^2,$$
$$\mathcal{O}_3 = i\vec{S}_N \cdot \left(\frac{\vec{q}}{m_N} \times \vec{v}^\perp\right),$$
$$\mathcal{O}_4 = \vec{S}_\chi \cdot \vec{S}_N,$$
$$\mathcal{O}_5 = i\vec{S}_\chi \cdot \left(\frac{\vec{q}}{m_N} \times \vec{v}^\perp\right),$$
$$\mathcal{O}_6 = \left(\vec{S}_\chi \cdot \frac{\vec{q}}{m_N}\right)\left(\vec{S}_N \cdot \frac{\vec{q}}{m_N}\right)$$
$$\mathcal{O}_7 = \vec{S}_N \cdot \vec{v}^\perp,$$
$$\mathcal{O}_8 = \vec{S}_\chi \cdot \vec{v}^\perp,$$
$$\mathcal{O}_9 = i\vec{S}_\chi \cdot \left(\vec{S}_N \times \frac{\vec{q}}{m_N}\right),$$
$$\mathcal{O}_{10} = i\vec{S}_N \cdot \frac{\vec{q}}{m_N},$$
$$\mathcal{O}_{11} = i\vec{S}_\chi \cdot \frac{\vec{q}}{m_N}.$$

… and much more considering experimental and theoretical uncertainties

DMp with preferred inelastic interaction:
$\chi^- + N \rightarrow \chi^+ + N$

- iDM mass states χ^+, χ^- with δ mass splitting
- Kinematic constraint for iDM:

$$\frac{1}{2}\mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

DAMA/NaI+DAMA/LIBRA
Slices from the 3d allowed volume in given scenario



Fund. Phys. 40(2010)900

iDM interaction on Tl nuclei of the NaI(Tl) dopant?
PRL106(2011)011301

- For large splittings, the dominant scattering in NaI(Tl) can occur off of Thallium nuclei, with A~205, which are present as a dopant at the 10^{-3} level in NaI(Tl) crystals.
- large splittings do not give rise to sizeable contribution on Na, I, Ge, Xe, Ca, O, … nuclei.

Mirror Dark Matter

Asymmetric mirror matter: mirror parity spontaneously broken ⇒ mirror sector becomes a heavier and deformed copy of ordinary sector
(See EPJC75(2015)400)

- Interaction portal: photon - mirror photon kinetic mixing $\frac{\epsilon}{2}F^{\mu\nu}F'_{\mu\nu}$
- mirror atom scattering of the ordinary target nuclei in the NaI(Tl) detectors of DAMA/LIBRA set-up with the Rutherford-like cross sections.

$\sqrt{f}\cdot\epsilon$ coupling const. and fraction of mirror atom



Perspectives for the future

Other signatures?

- *Diurnal effects*
- *Second order effects*
- *Shadow effects*
- *Directionality*
- *…*

Diurnal effects

EPJC 74 (2014) 2827

A diurnal effect with the sidereal time is expected for DM because of Earth rotation

Velocity of the detector in the terrestrial laboratory: $\vec{v}_{lab}(t) = \vec{v}_{LSR} + \vec{v}_{\odot} + \vec{v}_{rev}(t) + \vec{v}_{rot}(t),$

Since:

- $|\vec{v}_s| = |\vec{v}_{LSR} + \vec{v}_{\odot}| \approx 232 \pm 50$ km/s,
- $|\vec{v}_{rev}(t)| \approx 30$ km/s
- $|\vec{v}_{rot}(t)| \approx 0.34$ km/s at LNGS

$$v_{lab}(t) \simeq v_s + \hat{v}_s \cdot \vec{v}_{rev}(t) + \hat{v}_s \cdot \vec{v}_{rot}(t).$$

$\vec{v}_{LSR}$	velocity of the Local Standard of Rest (LSR) due to the rotation of the Galaxy
$\vec{v}_{\odot}$	Sun peculiar velocity with respect to LSR
$\vec{v}_{rev}(t)$	velocity of the revolution of the Earth around the Sun
$\vec{v}_{rot}(t)$	velocity of the rotation of the Earth around its axis at the latitude and longitude of the laboratory.

Annual modulation term:

$$\hat{v}_s \cdot \vec{v}_{rev}(t) = V_{Earth} B_m \cos(\omega(t - t_0))$$

- V_{Earth} is the orbital velocity of the Earth ≈ 30 km/s
- $B_m \approx 0.489$
- $t_0 \approx t_{equinox}$ + 73.25 days ≈ June 2

Diurnal modulation term:

$$\hat{v}_s \cdot \vec{v}_{rot}(t) = V_r B_d \cos\left[\omega_{rot}\,(t - t_d)\right]$$

- V_r is the rotational velocity of the Earth at the given latitude (for LNGS ≈ 0.3435 km/s)
- $B_d \approx 0.671$
- $t_d \approx 14.02$ h (at LNGS)



Earth velocity in the galactic frame. Starting time is spring equinox. The contribution of diurnal rotation has been dropped off. The maximum of the velocity (vertical line) is about 73 days after the spring equinox.



$v_s + v_s \cdot v_{rot}(t)$. Maximum of velocity about 14 h (vertical line).

Expected signal counting rate in a k−th energy bin:

$$S_k\left[v_{lab}(t)\right] \simeq S_k\left[v_s\right] + \left[\frac{\partial S_k}{\partial v_{lab}}\right]_{v_s} \left[V_{Earth} B_m \cos\omega(t - t_0) + V_r B_d \cos\omega_{rot}\,(t - t_d)\right]$$

The ratio R_{dy} is a model independent constant:

$$R_{dy} = \frac{S_d}{S_m} = \frac{V_r B_d}{V_{Earth} B_m} \simeq 0.016$$ at LNGS latitude

Diurnal effects in DAMA/LIBRA-phase1

EPJC 74 (2014) 2827



- Experimental *single-hit* residuals rate vs either sidereal and solar time.
- These residual rates are calculated from the measured rate of the *single-hit* events after subtracting the constant part

Energy	Solar Time	Sidereal Time
2–4 keV	$\chi^2/\text{d.o.f.} = 35.2/24 \rightarrow \text{P} = 7\%$	$\chi^2/\text{d.o.f.} = 28.7/24 \rightarrow \text{P} = 23\%$
2–5 keV	$\chi^2/\text{d.o.f.} = 35.5/24 \rightarrow \text{P} = 6\%$	$\chi^2/\text{d.o.f.} = 24.0/24 \rightarrow \text{P} = 46\%$
2–6 keV	$\chi^2/\text{d.o.f.} = 25.8/24 \rightarrow \text{P} = 36\%$	$\chi^2/\text{d.o.f.} = 21.2/24 \rightarrow \text{P} = 63\%$
6–14 keV	$\chi^2/\text{d.o.f.} = 25.5/24 \rightarrow \text{P} = 38\%$	$\chi^2/\text{d.o.f.} = 35.9/24 \rightarrow \text{P} = 6\%$

no diurnal variation with a significance of 95% C.L.

+ run test. The lower tail probabilities (in the four energy regions) are: 43, 18, 7, 26% for the solar case and 54, 84, 78, 16% for the sidereal case.

Thus, the presence of any significant diurnal variation and of time structures can be excluded at the reached level of sensitivity.

- Observed annual modulation amplitude in DAMA/LIBRA–phase1 in the (2–6) keV energy interval: (0.0097 ± 0.0013) cpd/kg/keV
- Thus, the expected value of the diurnal modulation amplitude is $\simeq 1.5 \times 10^{-4}$ cpd/kg/keV.
- When fitting the *single-hit* residuals with a cosine function with period fixed at 24 h and phase at 14 h: all the diurnal modulation amplitudes A_d are compatible with zero at the present level of sensitivity.

sidereal

Energy	A_d^{exp} (cpd/kg/keV)	χ^2/d.o.f.	P
2–4 keV	$(2.0 \pm 2.1) \times 10^{-3}$	27.8/23	22%
2–5 keV	$-(1.4 \pm 1.6) \times 10^{-3}$	23.2/23	45%
2–6 keV	$-(1.0 \pm 1.3) \times 10^{-3}$	20.6/23	61%
6–14 keV	$(5.0 \pm 7.5) \times 10^{-4}$	35.4/23	5%

A_d (2-6 keV) < 1.2×10^{-3} cpd/kg/keV (90%CL)

Present experimental sensitivity is not yet enough for the expected diurnal modulation amplitude derived from the DAMA/LIBRA–phase1 observed effect.

larger exposure DAMA/LIBRA–phase2 (+lower energy threshold) offers increased sensitivity to such an effect

Other signatures?

- *Diurnal effects*
- *Second order effects*
- *Shadow effects*
- *Directionality*
- *…*

Earth shadowing effect with DAMA/LIBRA–phase1



EPJC75(2015)239

- **Earth Shadow Effect** could be expected for DM candidate particles inducing nuclear recoils
- can be pointed out only for candidates with high cross-section with ordinary matter (low DM local density)
- would be induced by the variation during the day of the Earth thickness crossed by the DM particle in order to reach the experimental set-up



- DM particles crossing Earth lose their energy
- DM velocity distribution observed in the laboratory frame is modified as function of time (**GMST 8:00 black**; **GMST 20:00 red**)





Taking into account the DAMA/LIBRA DM annual modulation result, allowed regions in the ξ vs σ_n plane for each m_{DM}.

Investigation of Earth Shadow Effect

EPJC75 (2015) 239

Expected counting rate for a given mass, cross section and scenario by MC:

$$S_{d,sh}(t) = \xi\sigma_n S'_{d,sh}(t)$$

Expectations compared with diurnal residual rate of the *single-hit* events of DAMA/LIBRA-phase1 in (2-4) keV

Minimizing χ^2, upper limits on ξ can be evaluated





Considering DAMA/LIBRA DM annual modulation result, allowed regions in the ξ vs σ_n plane for each m_{DM}.

In these examples:
Isothermal halo model with v_0=220 km/s and v_{esc}=650 km/s

a) QF const. without channeling
b) QF const. including channeling
c) QF depending on energy
d) QF depending on energy renormalized to DAMA/LIBRA values

Red surface: 95% C.L. allowed mean value for ξ (surface thickness ± 30%)

DAMA/LIBRA phase2 - running

Quantum Efficiency features





Residual Contamination

The limits are at 90% C.L.

PMT	Time (s)	Mass (kg)	^{226}Ra (Bq/kg)	^{234m}Pa (Bq/kg)	^{235}U (mBq/kg)	^{228}Ra (Bq/kg)	^{228}Th (mBq/kg)	^{40}K (Bq/kg)	^{137}Cs (mBq/kg)	^{60}Co (mBq/kg)
Average			*0.43*	-	*47*	*0.12*	*83*	*0.54*	-	-
Standard deviation			*0.06*	-	*10*	*0,02*	*17*	*0.16*	-	-

Energy (keV)

Energy resolution



σ/E @ 59.5 keV for each detector with new PMTs with higher quantum efficiency (blu points) and with previous PMT EMI-Electron Tube (red points).

The light responses

Previous PMTs:	5.5-7.5 ph.e./keV
New PMTs:	up to 10 ph.e./keV

- To study the nature of the particles and features of related astrophysical, nuclear and particle physics aspects, and to investigate second order effects
- Special data taking for *other rare processes*

Features of the DM signal

The importance of studying second order effects and the annual modulation phase

High exposure and lower energy threshold can allow further investigation on:

- the nature of the DM candidates
- possible diurnal effects on the sidereal time
- astrophysical models

DAMA/NaI+LIBRA-phase1



The annual modulation phase depends on :

- Presence of **streams** (as SagDEG and Canis Major) in the Galaxy
- Presence of **caustics**
- Effects of gravitational **focusing of the Sun**

PRL112(2014)011301





The effect of the streams on the phase depends on the galactic halo model



A step towards such investigations:

DAMA/LIBRA-phase2 running with lower energy threshold

+ further possible improvements (DAMA/LIBRA-phase3) and DAMA/1ton

Possible DAMA/LIBRA-phase3

- The light collection of the detectors can further be improved
- Light yields and the energy thresholds will improve accordingly

The strong interest in the low energy range suggests the possibility of a new development of **high Q.E. PMTs** with **increased radiopurity** to directly couple them to the DAMA/LIBRA crystals, **removing** the special radio-pure quartz (Suprasil B) light guides (10 cm long), which act also as optical window.

The presently-reached PMTs features, but not for the same PMT mod.:

- Q.E. around 35-40% @ 420 nm (NaI(Tl) light)
- radiopurity at level of 5 mBq/PMT (^{40}K), 3-4 mBq/PMT (^{232}Th), 3-4 mBq/PMT (^{238}U), 1 mBq/PMT (^{226}Ra), 2 mBq/PMT (^{60}Co).

R&D efforts to obtain PMTs matching the best performances… feasible

No longer need for light guides (a 30-40% improvement in the light collection is expected)

… and multi-purpose DAMA/1ton

1) Proposed since 1996 (DAMA/NaI and DAMA/LIBRA intermediate steps)
2) Technology largely at hand and still room for further improvements in the low-background characteristics of the set-up (NaI(Tl) crystals, PMTs, shields, etc.)
3) 1 ton detector: the cheapest, the highest duty cycle, the clear signature, fast realization in few years

Design: DAMA/1ton can be realized by adding 3 replicas of DAMA/LIBRA:



- the detectors of similar size than those already used
- the features of low-radioactivity of the set-up and of all the used materials would be assured by many years of experience in the field
- electronic chain and controls would profit by the previous experience and by the use of compact devices already developped, tested and used.
- new digitizers will offer high expandibility and high performances
- the daq can be a replica of that of DAMA/LIBRA

• Some R&Ds carried out

Other signatures?

- *Diurnal effects*
- *Second order effects*
- *Shadow effects*
- *Directionality*
- *…*

THE DIRECTIONALITY APPROACH

(Approach that holds only for those DM candidates able to induce just nuclear recoils)

Based on the study of the correlation between the Earth motion in the galactic rest frame and the arrival direction of those DM candidates able to induce just nuclear recoils.



The dynamics of the rotation of the Milky Way galactic disc through the halo of DM causes the Earth to experience a wind of DM particles apparently flowing along a direction opposite to that of solar motion relative to the DM halo



… but because of the Earth's rotation around its axis, the DM particles average direction with respect to an observer fixed on the Earth changes during the sidereal day

The **direction of the induced nuclear recoils** can offer a way for pointing out the presence of those candidate particles; in fact the nuclear recoils are expected to be **strongly correlated** with their **impinging direction**, while the background events are not

DIRECTIONALITY SENSITIVE DETECTORS: ANISOTROPIC SCINTILLATORS

Anisotropic Scintillator:

- for heavy particles the light output and the pulse shape depends on the particle impinging direction with respect to the crystal axes
- for γ/e the light output and the pulse shape are isotropic



The variation of the response of an **anisotropic scintillator** during sidereal day can allow to point out the presence of a DM signal due to candidate inducing nuclear recoils

- The use of anisotropic scintillators to study the directionality signature was proposed for the first time in refs. [P. Belli et al., Il Nuovo Cim. C 15 (1992) 475; R. Bernabei et al., EPJC28(2003)203], where the case of anthracene was analysed; some preliminary activities have been carried out [N.J.C. Spooner et al, IDM1997 Workshop; Y. Shimizu et al., NIMA496(2003)347]

$ZnWO_4$ CRYSTAL SCINTILLATORS

- Low background $ZnWO_4$ crystal scintillators with large volume and good scintillation properties realized
- Various detectors with mass **0.1 - 0.7 kg** realized by exploiting different materials and techniques
- Detectors installed in a cavity (filled up with high-pure silicon oil) ϕ 47 x 59 mm in central part of a polystyrene light-guide 66 mm in diameter and 312 mm in length. The light-guides was faced by 2 low-background PMTs

PLB658(2008)193, NPA826(2009)256
NIMA626-627(2011)31,
JP38(2011)115107

Crystal scintillator	Size (mm)	Mass (g)
ZWO-1	20 × 19 × 40	117
ZWO-2	⌀44 × 55	699
ZWO-2a	⌀44 × 14	168





- Main aim of the measurements was the study of the properties of $ZnWO_4$ and the search for 2β processes in Zinc and Tungsten isotopes.





Advantages of the ZnWO4 crystal

- Very good anisotropic features
- High level of radiopurity
- High light output, that is low energy threshold feasible
- High stability in the running conditions
- Sensitivity to small and large mass DM candidate particles
- Detectors with ~ kg masses
- PSD capability



Light yield and energy threshold



Density (g/cm^3)	*7.87*
Melting point (°C)	*1200*
Structural type	*Wolframite*
Cleavage plane	*Marked (010)*
Hardness (Mohs)	*4–4.5*
Wavelength of emission maximum (nm)	*480*
Refractive index	*2.1–2.2*
Effective average decay time (μs)	*24*

PERFORMANCES OF $ZnWO_4$ CRYSTAL SCINTILLAOR

NIMA 626(2011)31

➢ *Radiopurity*

The measured radioactive contamination of $ZnWO_4$ approaches that of specially developed low background NaI(Tl):

- ~ 0.5 ppt for ^{232}Th;
- ~ 0.2 ppt for ^{238}U;
- < 0.02 mBq/kg for ^{40}K;
- total α activity of 0.18 mBq/kg



Run	Crystal	Size mass producer	t (h)	FWHM (%)	Ba
					0.2
1	ZWO-1	20 × 19 × 40 mm 117 g ISMA [a]	2906	12.6	1.7
2	ZWO-2	∅44 × 55 mm 699 g ISMA	2130	14.6	1.0
3	ZWO-3	∅27 × 33 mm 141 g ISMA (re-crystallization of ZWO-2)	994	18.2	1.5
4	ZWO-4	∅41 × 27 mm	834	14.2	2.3
				13.3	1.0



➢ *Pulse shape analysis*

The dependence of the pulse shapes on the type of irradiation in the $ZnWO_4$ scintillator allows one to discriminate $\beta(\gamma)$ events from those induced by α particles and to identify the α background

Developments is still ongoing: $\Rightarrow$ future $ZnWO_4$ crystals with higher radiopurity expected

ANISOTROPIC FEATURES IN $ZnWO_4$

The reachable sensitivity of the directionality approach depend on the anisotropic features of the detectors in response to the low energy nuclear recoils induced by the DM particles



Measurements with α **particles** have shown that the **light response** and the **pulse shape** of a $ZnWO_4$ depend on the impinging direction of α particles with respect to the crystal axes

Such effects are absent in case of electron excitation



These anisotropic effects are ascribed to preferred directions of the excitons' propagation in the crystal lattice affecting the dynamics of the scintillation mechanism

Similar effect is expected in the case of low energy nuclear recoils ⇒ Dedicated measurements are in preparation

Both the anisotropic features of the $ZnWO_4$ detectors can provide two independent ways to exploit the directionality approach

$ZnWO_4$ – work in progress…

At present:

- building a new dedicated experimental-setup for rare events at LNGS underground laboratory
- tests for light response a low temperature (~ -20 °C) of $ZnWO_4$ crystal scintillator
- tests about operational stability a low temperature (~ -20 °C) of $ZnWO_4$ crystal scintillator
- studies about light response vs neutron interactions in the $ZnWO_4$
- measurements about new technique in order to develop $ZnWO_4$ crystal scintillator with an extremely high level of radiopurity





EXAMPLE OF THE EXPECTED SIGNAL IN A SIMPLIFIED MODEL CONSIDERED IN EPJC73(2013)2276

Expected signal rate as a function of sidereal time and days of the year

[2-3] keV $\sigma_p=5\times10^{-5}$ pb
m_{DM}= 10 GeV



[6-7] keV $\sigma_p=5\times10^{-5}$ pb
m_{DM}= 100 GeV



Eur. Phys. J. C 73 (2013) 2276

MODEL DEPENDENT COMPARISONS; EXAMPLE OF REACHABLE SENSITIVITY IN A SCENARIO CONSIDERED IN EPJC73(2013)2276

Considering an experiment with:

- **200 kg of $ZnWO_4$;**
- **5 years of data taking.**

The reachable sensitivity has been calculated considering four possible time independent **background levels** in the low energy region:

- **10^{-4} cpd/kg/keV** ——————
- **10^{-3} cpd/kg/keV** – – – – –
- **10^{-2} cpd/kg/keV** .....
- **0.1 cpd/kg/keV** — · — · —

Black lines are the **sensitivities reachable** with four possible background levels in the low energy region in a given scenario

Eur. Phys. J. C 73 (2013) 2276





The directionality approach can reach in the given scenario a sensitivity to the cross section at level of 10^{-5} – 10^{-7} pb, depending on the particle mass

For comparison, there are also shown (green, red and blue) allowed regions obtained with a corollary analysis of the 9.3 σ C.L. DAMA/NaI + DAMA/LIBRA model independent result in terms of scenarios for the DM candidates considered here

Phys. Rev. D 84, 055014 (2011)

Conclusions

- Positive evidence for the presence of DM particles in the galactic halo supported at **9.3σ** C.L. (14 annual cycles DAMA/NaI and DAMA/LIBRA-phase1: 1.33 ton × yr)
- Modulation parameters determined with high precision
- New investigation on different peculiarities of the DM signal exploited (**Diurnal Modulation** and **Earth Shadow Effect**)
- Full sensitivity to many kinds of DM candidates and interactions types (both inducing recoils and/or e.m. radiation), full sensitivity to low and high mass candidates

- **DAMA/LIBRA – phase2 in data taking** at lower software energy threshold (below 2 keV) to investigate further features of DM signals and second order effects
- Continuing investigations of rare processes other than DM as well as further developments
- **DAMA/LIBRA – phase3 R&D in progress**
- R&D for a possible DAMA/1ton set-up, proposed by DAMA since 1996, **continuing**
- Study of $ZnWO_4$ scintillator for exploiting directionality technique **in progress**

Thank you for your attention