




*symmetry*

an open access journal by 

# Maxim Yu. Khlopov

VIA, APC Laboratory, Paris, France;

National Research Nuclear University “MEPHI”

(Moscow Engineering and Physics Institute) Moscow, Russia

and

Institute of Physics, Southern Federal University, Rostov on Don, Russia

## **Conspiracy of BSM physics and cosmology**

Talk at

**XXII Bled Workshop**

**“What comes beyond the Standard models?”**

Bled, Slovenia 9 July 2019

*Sponsored by « Symmetry »*

# Outlines

- Arguments for BSM Physics
- Mass of neutrino – the only proven BSM phenomenon
- BSM Physical basis of the SM cosmology
- BSM cosmology that follows from BSM physics.
- Primordial Black Holes (PBHs) as cosmological reflection of particle symmetry.
- Strong Primordial nonhomogeneities and Massive PBH clusters from models of inflation.
- Antimatter as profound signature for nonhomogeneous baryosynthesis.

# The Standard Model (SM) of Particle Physics

- Gauge principle (Quantum Electrodynamics (QED)):


$$L = i\bar{\psi}\hat{\partial}\psi + eA_\mu J_\mu - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}, \quad J_\mu = \bar{\psi}\gamma_\mu\psi, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

$$\psi \rightarrow \psi \exp(ie\chi(x)) \quad \longleftarrow \quad U(1)\text{-symmetry transformation}$$

$$A_\mu \rightarrow A_\mu + \partial_\mu\chi$$

- Generalization on *weak* and *strong* interactions:  $SU(3)_c \times SU(2)_L \times U(1)_Y = \mathbf{SM}$
- *Higgs mechanism* of spontaneous symmetry breaking

$$m_f = \frac{fv}{\sqrt{2}}, \quad m_W = \frac{fv}{2}, \quad m_H = \sqrt{2\lambda}v$$



# Problems of SM (1)

$$SU(3)_c \times SU(2)_L \times U(1)_Y$$

1) Excessive number of *parameters*:  $g, g', v, \lambda, \{f_{ij}\}, \dots$

2) Problem of *number of generations*

$$\begin{pmatrix} \nu_e \\ e \end{pmatrix} \begin{pmatrix} \nu_\mu \\ \mu \end{pmatrix} \begin{pmatrix} \nu_\tau \\ \tau \end{pmatrix}$$
$$\begin{pmatrix} u \\ d \end{pmatrix} \begin{pmatrix} c \\ s \end{pmatrix} \begin{pmatrix} t \\ b \end{pmatrix}$$

3) The problem of *neutrino mass*.

$$m(\nu_e) < 1 \text{ eV} \ll m(e) = 511 \text{ keV}$$

4) *CKM* mixing matrix  $\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} & & \\ & 3 & \\ & & 3 \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$

5) The problem of *C-, P- and CP-violation* observed (in weak interactions and K-mesons).

# Problems of SM (2)



$$SU(3)_c \times SU(2)_L \times U(1)_Y$$

## 6) Problem of *CP-violation in QCD*

$$\Delta L_{\text{QCD}} = \frac{\alpha_s^2}{16\pi} \theta \cdot \varepsilon_{\alpha\beta\mu\nu} G^{a\alpha\beta} G^{a\mu\nu} \quad \longrightarrow \quad \text{Strong CP-violation}$$

$$\longrightarrow \quad d_n \approx e\theta m_u / m_n^2 \sim \theta \cdot 10^{-16} e \cdot \text{cm} \quad \text{From experiment one has } |\theta| < 10^{-9}$$

## 7) Divergence of *Higgs boson mass* (due to radiation corrections), and, as consequence, instability of Higgs potential against radiation corrections



# Other arguments for extension of SM


## $SU(3)_c \times SU(2)_L \times U(1)_Y$

*Unifying all interactions:*

*for aesthetical reasons*

*for reasons of similarity in their descriptions:  $L \sim g A_\mu J_\mu$*

*convergence of gauge constants (SUSY changes the lines and gives it naturally)*



# Trends in extension of SM (1)

“See-saw” mechanism of generation of neutrino mass

$$\Delta L_m = m_D (\bar{\nu}_L \nu_R + \bar{\nu}_R \nu_L) + m_R (\bar{\nu}_R^c \nu_R + \bar{\nu}_R \nu_R^c) + m_L (\bar{\nu}_L^c \nu_L + \bar{\nu}_L \nu_L^c)$$

To avoid problems with SM parameters

at  $m_D \ll m_R$


$$\Delta L_m = m_1 (\bar{\nu}_1^c \nu_1 + \bar{\nu}_1 \nu_1^c) + m_2 (\bar{\nu}_2^c \nu_2 + \bar{\nu}_2 \nu_2^c)$$

$$m_1 \approx \frac{m_D^2}{m_R} \ll m_D \sim m_l, \quad m_2 \approx m_R$$

Majorana mass term violates lepton number, which will lead to processes with  $\Delta L=2$ .

Consequence: neutrinoless double  $\beta$ -decay


Spontaneous breaking of symmetry of lepton charge implies the existence of Nambu-Goldstone boson – Majoron.



# Trends in extension of SM (2)

## “Mirror” world

Kobzarev, Okun, Pomeranchuk (1966)



*The equivalence between left- and right-handed coordinate systems is restored, if reflection in mirror is accompanied by change of ordinary particles by their mirror partners. Mirror partners are strictly symmetric to ordinary particles. Therefore they can not have ordinary electromagnetic and strong interactions (doubling of atomic levels, or pion states). Successive analysis have shown that (O) and (M) also can not share W and Z boson mediated weak interaction.* 8



# Trends in extension of SM (3)

## *SU(5) GUT model*

$$SU(5) \supset SU(3)_c \times SU(2)_L \times U(1)_Y$$

- There are **24 gauge bosons**, which contain 12 known ones ( $g, Z, W, \gamma$ ) and 12 new ones ( $X, Y$ )
- All known fermions (15, excluding  $\nu_R$ ) are distributed into  $SU(5)$  5-plets and 10-plets.
- $\Lambda_{GUT} \sim 10^{15} \text{ GeV}$

Example of 5-plet:  $(\bar{d}_{red} \quad \bar{d}_{yellow} \quad \bar{d}_{blue} \quad e^- \quad \nu_e)_L$

Proton decays


Note:  $\Delta B = \Delta L$

$$\tau_p^{\text{theor}} \sim \frac{\Lambda_{GUT}^5}{m_p^4} \sim 10^{31} \text{ years}$$

vs

$$\tau_{p \rightarrow e\pi}^{\text{exp}} > 10^{33} \text{ years}$$

*It strongly restricts the model.*



# Trends in extension of SM (4)

## Symmetry of families and Familons

To resolve **the problem of quark-lepton families** a new global symmetry  $SU(3)_H$  can be introduced. It's spontaneous breaking leads to 8 new scalar fields Nambu-Goldstone bosons - **familons**.

Familons interact with quarks and leptons at energy scale  $v_a$  as

$$L_{aff} = ic_{aff} \frac{m_f}{v_a} \bar{a}f \gamma_5 f$$

Their interactions lead to various transitions between fermions of different families.

In the model of singlet familon there is only 1 such Nambu-Goldstone boson.

# Trends in extension of SM (5)

## Model of Peccei-Quinn

To resolve **the problem of strong CP-violation** a new global symmetry  $U(1)_{PQ}$  with new scalar field is introduced.

Scalar field is associated with spontaneous  $U(1)_{PQ}$  violation at energy scale  $v_a$ , and provides mechanism of dynamical suppression of  $\theta$  (due to its Nambu-Goldstone boson – **axion**).

Axion interacts with quarks as

$$L_{aff} = ic_{aff} \frac{m_f}{v_a} a \bar{f} \gamma_5 f$$

and have potential  $V_{eff} \sim m_u \Lambda_{QCD}^3 \left(1 - \cos(\theta + a / v_a)\right)$

minimizing at  $a = \langle a \rangle = -\theta v_a$

$$\longrightarrow \Delta L_{QCD} = \frac{\alpha_s^2}{16\pi} \left( \theta + \frac{a}{v_a} \right) \cdot \epsilon_{\alpha\beta\mu\nu} G^{a\alpha\beta} G^{a\mu\nu} \rightarrow 0$$

# Trends in extension of SM (5)

## Model of Peccei-Quinn: properties of axion


The axion has the mass, which is related with its energy scale

$$m_a \approx 0,6 \text{ meV} \frac{10^{10} \text{ GeV}}{v_a}$$

hep-ph/0002154

It decays into two photons at


$$\begin{aligned} \tau(a \rightarrow \gamma\gamma) &= \frac{64\pi^3}{c_{a\gamma\gamma}^2 \alpha^2} \frac{v_a^2}{m_a^3} \approx \frac{2 \cdot 10^{17} \text{ years}}{c_{a\gamma\gamma}^2} \left( \frac{\text{eV}}{m_a} \right)^5 \approx \\ &\approx \frac{2 \cdot 10^{33} \text{ years}}{c_{a\gamma\gamma}^2} \left( \frac{v_a}{10^{10} \text{ GeV}} \right)^5. \end{aligned}$$



According existing constraints its mass and energy scale must be

$$0.5 \cdot 10^{-5} \text{ eV} < m_a < 0,5 \cdot 10^{-2} \text{ eV}$$

$$2 \cdot 10^9 \text{ GeV} < v_a < 10^{12} \text{ GeV}$$



# Trends in extension of SM (6)

## Supersymmetry (SUSY)

All particles with spin  $s$  get supersymmetric partner – particle with spin  $s' = |s - 1/2|$  and all other identical quantum numbers (except the mass in case of broken SUSY).

$$e_{L,R}(s=0) \leftrightarrow \tilde{e}_{L,R}(s=0)$$


$$\gamma(s=1) \leftrightarrow \tilde{\gamma}(s=1/2)$$

$$\text{or } B(s=1) \leftrightarrow \tilde{B}(s=1/2)$$

....

Photino (bino) is Majorana particle

1) **SUSY** helps to resolve the problem of large radiation correction in Higgs sector.



2) **SUSY** improves convergence of the gauge constants (in GUT).


3) **SUSY** may serve as a basic symmetry for construction of quantum gravity.

**SUSY must be, evidently, broken.** A “hidden” sector is introduced which induces soft SUSY breaking in our sector by means of 1) gravity (**SUGRA**), 2) special gauge fields (**GMSB**). 13

# Trends in extension of SM (6)

## Supersymmetry (SUSY)

To avoid fast decay of *proton* in GUT, a multiplicative quantum number – *R-parity* is introduced.  $R=+1$  is assigned for all ordinary particles, and  $R=-1$  – for SUSY partners.



*Consequence:* the lightest SUSY particles (**LSP**) is stable.

**Neutralino** is often suggested for a role of LSP. Neutralino is a superposition of four SUSY partners of bino, neutral wino, two neutral higgsino.

$$\chi_{1,2,3,4}^0 = N_{1,2,3,4}^{(B)} \tilde{B} + N_{1,2,3,4}^{(W)} \tilde{W}^3 + N_{1,2,3,4}^{(H_1)} \tilde{H}_1^0 + N_{1,2,3,4}^{(H_2)} \tilde{H}_2^0$$

Also, **gravitino** or **axino** are sometimes suggested for a role of LSP.

# Trends in extension of SM (7)


## Kaluza-Klein (KK) models

$$g_{MN} = \begin{pmatrix} g_{\mu\nu} & g_{\mu n} \\ g_{m\nu} & g_{mn} \end{pmatrix}$$


*Describes other interactions*

$M, N = 0, \dots, D$   
 $\mu, \nu = 0, 1, \dots, 3$   
 $m, n = 4, \dots, D$   
 $\underbrace{\hspace{10em}}_d$

If extra  $d$  dimensions are compactified with  $R = \text{const}$ , then they are manifested in form of the *KK-mass states* for all particles, if  $R \neq \text{const}$  – in form of *interaction*.



# Composition of the Modern Universe



$$\Omega \equiv \frac{\rho}{\rho_{cr}}$$

$$\Omega_b \approx 0.044 \quad \Omega_{\text{CMB}} \approx 0.5 \cdot 10^{-4}$$

$$\Omega_{\text{DM}} \approx 0.20$$

$$\Omega_{\Lambda} \approx 0.7$$

$$\Omega_{\text{tot}} \approx 1.0$$

**In the modern Universe dominate dark energy and dark matter – their nature is related to the new physics – physics beyond the Standard model, on which the bedrocks of modern cosmology are based**



# The bedrocks of modern cosmology

*Our current understanding of structure and evolution of the Universe implies three necessary elements of Big Bang cosmology that can not find physical grounds in the standard model of electroweak and strong interactions. They are:*


- Inflation
- Baryosynthesis
- Dark matter/energy

*Physics beyond the Standard model, describing these phenomena inevitably predicts additional model dependent effects, in which NS and BH play important role.*

# Cosmological Reflections of Microworld Structure

- **(Meta-)stability of new particles reflects some Conservation Law, which prohibits their rapid decay. Following Noether's theorem this Conservation Law should correspond to a (nearly) strict symmetry of microworld. Indeed, all the particles - candidates for DM reflect the extension of particle symmetry beyond the Standard Model.**
- **In the early Universe at high temperature particle symmetry was restored. Transition to phase of broken symmetry in the course of expansion is the source of topological defects (monopoles, strings, walls...).**
- **Structures, arising from dominance of superheavy metastable particles and phase transitions in early Universe, can give rise to Black Holes, retaining in the Universe after these structures decay.**

# Cosmological Dark Matter



***Cosmological Dark Matter explains:***

- ***virial paradox in galaxy clusters,***
- ***rotation curves of galaxies***
- ***dark halos of galaxies***
- ***effects of macro-lensing***

***But first of all it provides formation of galaxies from small density fluctuations, corresponding to the observed fluctuations of CMB***

*To fulfil these duties Dark Matter should interact sufficiently weakly with baryonic matter and radiation and it should be sufficiently stable on cosmological timescale.*

*Baryon density estimated from the results of BBN (mainly from Primordial deuterium) is not sufficient to explain the matter content of the modern Universe*

# **Dark Matter – Cosmological Reflection of Microworld Structure**

**Dark Matter should be present in the modern Universe, and thus is stable on cosmological scale.**

**This stability reflects some Conservation Law, which prohibits DM decay.**

**Following Noether's theorem this conservation law should correspond to a (nearly) strict symmetry of microworld.**

# Dark Matter from Elementary Particles

*By definition Dark Matter is non-luminous, while charged particles are the source of electromagnetic radiation. Therefore, neutral weakly interacting elementary particles are usually considered as Dark Matter candidates. If such neutral particles with mass  $m$  are stable, they freeze out in early Universe and form structure of inhomogeneities with the minimal characteristic scale*

$$M = m_{Pl} \left( \frac{m_{Pl}}{m} \right)^2$$

- However, if charged particles are heavy, stable and bound within neutral « atomic » states they can also play the role of specific composite Dark matter (Dark atoms).
- Physical models, underlying dark atom scenarios, their problems and nontrivial solutions as well as the possibilities for their test will be the subject of the successive talks.

**THE PUZZLES  
OF DIRECT DARK MATTER  
SEARCHES**

# Direct searches for Dark Matter

## **Possibility of detecting relict massive neutrinos**

V. F. Shvartsman, V. B. Braginskii, S. S. Gershtein, Ya. B. Zel'dovich, and  
M. Yu. Khlopov

*M. V. Keldysh Institute of Applied Mathematics, Academy of Sciences of the USSR*


(Submitted 18 August 1982)

*Pis'ma Zh. Eksp. Teor. Fiz.* **36**, No. 6, 224–226 (20 September 1982)

The coherent intensification of the interaction of relict massive neutrinos with grains of matter with a size on the order of the neutrino wavelength suggests that it might be possible to detect a galactic neutrino sea by virtue of the mechanical pressure which it exerts in the direction opposite that in which the solar system is moving in the galaxy.

# WIMP-nucleus interaction

CDM can consist of Weakly Interacting Massive Particles (WIMPs).  
Such particles can be searched by effects of WIMP-nucleus interactions.



$$\Delta T = 0 \div \Delta T_{\max} = \frac{q_{\max}^2}{2m_A} = \frac{2\mu^2 v^2}{m_A} \xrightarrow{m_X \gg m_A} 2m_A v^2$$

$$v \sim 300 \frac{\text{km}}{\text{s}}, \quad \Delta T \sim 10 \text{keV}$$

$$\Delta T_{\max} < E_A^* \Rightarrow \text{elastic scattering}$$

$$q_{(\max)} R_A > 1 \Rightarrow \text{non-pointlike nucleus}$$

$$\text{Interaction amplitude} \equiv A_{AX} = A_{AX}^{\text{point}} \cdot F_A(q^2)$$




# Annual modulation of WIMP effects

## Minimization of background

- Installation deeply underground
- Radioactively pure materials
- Annual modulation

DM does not participate in rotation around GC.



$$v_{Earth\ y} = v_{Sun\ y} + v_{orb} \cdot \cos \gamma \cdot \cos \omega(t - t_0)$$

$$v_{Sun\ y} = 220 + 16.5 \cdot \cos 25^\circ \cdot \sin 53^\circ \text{ (km/s)}$$

$$t_0 = 2 \text{ June}$$

$$Amplitude < \sim \frac{v_{orb} \cdot \cos \gamma}{v_{Sun\ y}} \sim \frac{15}{232} \sim 7\%$$

**THE PUZZLES  
OF INDIRECT DARK MATTER  
SEARCHES**

# Indirect searches for Dark Matter

## Astrophysical bounds on the mass of heavy stable neutral leptons

Ya. B. Zel'dovich, A. A. Klypin, M. Yu. Khlopov, and V. M. Chechetkin

*Institute of Applied Mathematics, USSR Academy of Sciences*

(Submitted 29 November 1979)

Yad. Fiz. **31**, 1286–1294 (May 1980)

Analytical and numerical calculations show that heavy neutral stable leptons are carried along by the collapsing matter during the formation of galaxies and possibly stars as well. The condensation in galaxies and stars results in appreciable annihilation of leptons and antileptons. Modern observations of cosmic-ray and  $\gamma$ -ray fluxes establish a limit  $m_\nu \gtrsim 100$  GeV for the mass of neutral leptons, since annihilation of neutral leptons produces  $\gamma$  rays and cosmic rays. The obtained bound, in conjunction with ones established earlier, precludes the existence of stable neutral leptons (neutrinos) with  $m_\nu > 30$  eV.

# Condensation of Dark Matter in Galaxy


$$\ddot{R} + \omega^2 R = 0$$

$$\omega^2 = 4\pi G(\rho_v + \rho_b)$$

$$I = \frac{E(t)}{\omega(t)} = \frac{\omega^2 R^2}{2\omega} = \text{const}$$

$$\rho_v(t) \propto R^{-3} \propto \omega^{3/2} \propto [\rho_b(t)]^{3/4}$$

$$\rho_v(t) \propto [\rho_b(t)]^{3/4}$$




- Motion of collisionless gas in nonstationary field of baryonic matter, contracting owing to dissipation processes, provides effective dissipation and contraction of this gas.
- In result collisionless Dark Matter condenses in Galaxy, but it is distributed more steeply, than baryonic matter.
- It qualitatively explains the difference in distribution of baryons and dark matter.
- Due to condensation effects of annihilation in Galaxy can be significant even for subdominant DM components (e.g.4th neutrino).

# Annihilation and decays of DM as a source of CR.

Stable DM particles can annihilate



$$\dot{n}_{sources} = n_X n_{\tilde{X}} \langle \sigma_{ann} v \rangle$$



Metastable neutral particles decay with equal amount of positrons and electrons



At the level of elementary process metastable double charged particles can decay to same sign leptons only



The excess of high energy positrons detected in PAMELA, FERMI/LAT and AMS02 experiments may be considered as an evidence for indirect effect of dark matter, first predicted by Zeldovich et al (1980).

# Cosmic positron excess from DM?





Figure 3: Positron excess due to  $UU \rightarrow e^+e^+, \mu^+\mu^+, \tau^+\tau^+$  decays compared to PAMELA and AMS-02 data.

Probably such indirect effect is detected in the cosmic positron fluxes.

[figure from K.M.Belotsky et al. arXiv:1403.1212]

# AMS02 in the next decade



*Presented in CERN on 08.12.2016 by Prof. S.Ting*

# INTEGRAL excess of positron annihilation line

- In the galactic bulge the excess of positron annihilation line is observed by INTEGRAL.
- This effect may be due to extra positrons originated from dark matter.



# **DARK MATTER FROM CHARGED PARTICLES?**

# Baryonic Matter – atoms of stable quarks and charged lepton (electron)

- Ordinary matter consists of atoms
- Atoms consist of nuclei and electrons.
- Electrons are lightest charged particles – their stability is protected by the conservation of electric charge.
- Nuclei consist of nucleons, whose stability reflects baryon charge conservation.

In ordinary matter stable elementary particles are electrically charged, but bound in neutral atoms.

# Dark Matter from Charged Particles?

*By definition Dark Matter is non-luminous, while charged particles are the source of electromagnetic radiation. Therefore, neutral weakly interacting elementary particles are usually considered as Dark Matter candidates. If such neutral particles with mass  $m$  are stable, they freeze out in early Universe and form structure of inhomogeneities with the minimal characteristic scale*

$$M = m_{Pl} \left( \frac{m_{Pl}}{m} \right)^2$$

- However, if charged particles are heavy, stable and bound within neutral « atomic » states they can play the role of composite Dark matter.
- Physical models, underlying such scenarios, their problems and nontrivial solutions as well as the possibilities for their test are the subject of the present talk.

# « No go theorem » for -1 charge components

- *If composite dark matter particles are « atoms », binding positive P and negative E charges, all the free primordial negative charges E bind with He-4, as soon as helium is created in SBBN.*
- *Particles E with electric charge -1 form +1 ion [E He].*
- *This ion is a form of anomalous hydrogen.*
- *Its Coulomb barrier prevents effective binding of positively charged particles P with E. These positively charged particles, bound with electrons, become atoms of anomalous isotopes*
- *Positively charged ion is not formed, if negatively charged particles E have electric charge -2.*

# Nuclear-interacting composite dark matter: O-helium « atoms »



If we have a stable double charged particle  $X^{--}$  in excess over its partner  $X^{++}$  it may create Helium like neutral atom (O-helium) at temperature  $T < I_0$ ,

Where:



${}^4\text{He}$  is formed at  $T \sim 100 \text{ keV}$  ( $t \sim 100 \text{ s}$ )

This means that it would rapidly create a neutral atom, in which all  $X^{--}$  are bound



The Bohr orbit of O-helium « atom » is of the order of radius of helium nucleus.



## References

1. M.Yu. Khlopov, *JETP Lett.* 83 (2006) 1;
2. D. Fargion, M.Khlopov, C.Stephan, *Class. Quantum Grav.* 23 (2006) 7305;
2. M. Y. Khlopov and C. Kouvaris, *Phys. Rev. D* 77 (2008) 065002]

# Constituents of composite dark matter

*Few possible candidates for -2 charges:*

*Stable doubly charged "leptons" with mass  $>100$  GeV ( $\sim 1$  TeV range):*

- *AC « leptons » from almost commutative geometry*

D. Fargion, M.Khlopov, C.Stephan, Class. Quantum Grav. 23 (2006) 7305

- *Technibaryons and technileptons from Walking Technicolor (WTC)*

M. Y. Khlopov and C. Kouvaris, Phys. Rev. D 77 (2008) 065002; M. Y. Khlopov and C. Kouvaris, Phys. Rev. D 78 (2008) 065040

*Hadron-like bound states of:*

- *Stable U-quark of 4-th family in Heterotic string phenomenology*

M.Yu. Khlopov, JETP Lett. 83 (2006) 1

- *Stable U-quarks of 5th family in the approach, unifying spins and charges*

N.S. Mankoc Borstnik, Mod. Phys. Lett. A 10 (1995) 587

M.Yu.Khlopov, A.G.Mayorov, E.Yu.Soldatov (2010), arXiv:1003.1144

# WTC-model

*The ideas of Technicolor (TC) are revived with the use of  $SU(2)$  group for “walking” (not running ) TC gauge constant \*.*

- 1.  $U$  and  $D$  techniquarks bound by Technicolor give mass to  $W$  and the  $Z$  bosons.*
- 2.  $UU$ ,  $UD$ ,  $DD$  and their corresponding antiparticles are technibaryons and corresponding anti-technibaryons.*
- 3. The electric charges of  $UU$ ,  $UD$ , and  $DD$  are in general  $y+1$ ,  $y$  and  $y-1$  respectively, where  $y$  is an arbitrary real number.*
- 4. In order to cancel the **Witten global anomaly** the model requires in addition an existence of a fourth family of leptons.*
- 5. Their electric charges are in terms of  $y$  respectively  $(1 - 3y)/2$  and  $(-1 - 3y)/2$ .  
If  $y=1$ , both **stable doubly charged** technibaryons and technileptons are possible\*\*.*

*All these stable techniparticles will look like stable multiple charged leptons at LHC*

## References

\*

F. Sannino and K. Tuominen, *Phys. Rev. D* 71 (2005) 051901 ;  
D. K. Hong *et al.*, *Phys. Lett. B* 597 (2004) 89 ;  
D. D. Dietrich *et al.*, *Phys. Rev. D* 72 (2005) 055001 ;  
S. B. Gudnason *et al.*, *Phys. Rev. D* 73 (2006) 115003 ;  
S. B. Gudnason *et al.*, *Phys. Rev. D* 74 (2006) 095008]

\*\*

M. Y. Khlopov and C. Kouvaris, *Phys. Rev. D* 77 (2008) 065002;

# Techniparticle excess

- The advantage of WTC framework is that it provides definite relationship between baryon asymmetry and techniparticle excess.

$$\frac{TB}{B} = -\sigma_{UU} \left( \frac{L}{B} \frac{1}{3\sigma_{\zeta}} + 1 + \frac{L}{3B} \right)$$


Here  $\sigma_i$  ( $i = UU, \zeta$ ) are statistical factors in equilibrium relationship between, TB, B, L and L'

*The equilibrium is maintained by electroweak SU(2) sphalerons and similar relationship can hold true for any SU(2) dublets (like U quarks of 4th family or stable quarks of 5th family)*




# Relationship between TB and B

$$\xi = \frac{L'}{3B\sigma_\zeta} + 1 + \frac{L}{3B}$$




- $L'=0, T^*=150$  GeV  
 $\xi = 0.1; 1; 4/3; 2; 3$



$$\xi = 4/3$$

- $L'=0,$   
 $T^*=150, 200, 250$  GeV

# Relationship between TB, L' and B



- $x$  denotes the fraction of dark matter given by the technibaryon
- $TB < 0, L' > 0$  – two types of -2 charged techniparticles.


*The case  $TB > 0, L' > 0$  ( $TB < 0, L' < 0$ ) gives an interesting possibility of (-2 +2) atom-like WIMPs, similar to AC model. For  $TB > L'$  ( $TB < L'$ ) no problem of free +2 charges*

# Stable multiple charged particles

*WTC can lead to techniparticles with multiple charge*

$q$	$UU(q + 1)$	$UD(q)$	$DD(q - 1)$	$\nu'(\frac{1-3q}{2})$	$\zeta(\frac{-1-3q}{2})$
1	2	1	0	-1	-2
3	4	3	2	-4	-5
5	6	5	4	-7	-8
7	8	7	6	-10	-11

*-2n charged particles in WTC bound with n nuclei of primordial He form Thomson atoms of XHe*



# **O-HELIUM DARK MATTER**

# O-helium dark matter

$$T < T_{od} = 1keV$$

$$n_b \langle \sigma v \rangle \left( m_p / m_o \right) t < 1$$

$$T_{RM} = 1eV$$

$$M_{od} = \frac{T_{RM}}{T_{od}} m_{Pl} \left( \frac{m_{Pl}}{T_{od}} \right)^2 = 10^9 M_{Sun}$$

- Energy and momentum transfer from baryons to O-helium is not effective and O-helium gas decouples from plasma and radiation
- O-helium dark matter starts to dominate
- On scales, smaller than this scale composite nature of O-helium results in suppression of density fluctuations, making O-helium gas Warmer than Cold Dark Matter

# O-helium in Earth

- Elastic scattering dominates in the (OHe)-nucleus interaction. After they fall down terrestrial surface the in-falling OHe particles are effectively slowed down due to elastic collisions with the matter. Then they drift, sinking down towards the center of the Earth with velocity

$$V = \frac{g}{n\sigma v} \approx 80S_3 A_{med}^{1/2} \text{ cm/ s.}$$

Here  $A_{med} \sim 30$  is the average atomic weight in terrestrial surface matter,  $n = 2.4 \cdot 10^{24}/A_{med}$  is the number of terrestrial atomic nuclei,  $\sigma v$  is the rate of nuclear collisions and  $g = 980 \text{ cm/ s}^2$ .

# O-helium experimental search?

- In underground detectors, (OHe) “atoms” are slowed down to thermal energies far below the threshold for direct dark matter detection. However, (OHe) nuclear reactions can result in observable effects.
- O-helium gives rise to less than 0.1 of expected background events in XQC experiment, thus avoiding severe constraints on Strongly Interacting Massive Particles (SIMPs), obtained from the results of this experiment.

*It implies development of specific strategy for direct experimental search for O-helium.*

# **O-HELIUM DARK MATTER IN UNDERGROUND DETECTORS**



# O-helium concentration in Earth

The O-helium abundance the Earth is determined by the equilibrium between the in-falling and down-drifting fluxes.

The in-falling O-helium flux from dark matter halo is

$$F = \frac{n_0}{8\pi} \cdot |\mathbf{V}_h + \mathbf{V}_E|,$$

where  $\mathbf{V}_h$  is velocity of Solar System relative to DM halo (220 km/s),  $\mathbf{V}_E$  is velocity of orbital motion of Earth (29.5 km/s) and

$n_0 = 3 \cdot 10^{-4} S_3^{-1} \text{ cm}^{-3}$  is the local density of O-helium dark matter.

At a depth  $L$  below the Earth's surface, the drift timescale is  $\sim L/V$ . It means that the change of the incoming flux, caused by the motion of the Earth along its orbit, should lead at the depth  $L \sim 10^5 \text{ cm}$  to the corresponding change in the equilibrium underground concentration of OHe on the timescale

$$t_{dr} \approx 2.5 \cdot 10^2 S_3^{-1} \text{ s}$$

# Annual modulation of O-helium concentration in Earth

The equilibrium concentration, which is established in the matter of underground detectors, is given by

$$n_{\text{oE}} = \frac{2\pi \cdot F}{V} = n_{\text{oE}}^{(1)} + n_{\text{oE}}^{(2)} \cdot \sin(\omega(t - t_0)),$$

where  $\omega = 2\pi/T$ ,  $T=1\text{yr}$  and  $t_0$  is the phase. The averaged concentration is given by



$$n_{\text{oE}}^{(1)} = \frac{n_0}{320S_3A_{\text{med}}^{1/2}}V_h$$

and the annual modulation of OHe concentration is characterized by


$$n_{\text{oE}}^{(2)} = \frac{n_0}{640S_3A_{\text{med}}^{1/2}}V_E$$

**The rate of nuclear reactions** of OHe with nuclei is proportional to the local concentration and the energy release in these reactions leads to ionization signal containing both constant part and **annual modulation**.


# OHe solution for puzzles of direct DM search

- OHe equilibrium concentration in the matter of DAMA detector is maintained for less than an hour 
- Annual modulations in inelastic processes, induced by OHe in matter. No signal of WIMP-like recoil
- The process   
 $OHe + (A, Z) \Rightarrow [OHe(A, Z)] + \gamma$   
is possible, in which only a few keV energy is released. Other inelastic processes are suppressed
- Signal in DAMA detector is not accompanied by processes with large energy release. This signal corresponds to a formation of anomalous isotopes with binding energy of few keV

# Potential of OHe-nucleus interaction




$$U_{Xnuc} = -2Z\alpha \left( \frac{1}{r} + \frac{1}{r_0} \right) \exp(-2r/r_0)$$



$$U_{Stark} = -\frac{2Z\alpha}{r^4} \frac{9}{2} r_0^3$$




$$U_{Coul} = +\frac{2\alpha Z}{\rho} - \frac{2\alpha Z}{r}$$



$$U_{osc} = -\left[ \frac{(Z+2)\alpha}{R} \left( 1 - \left( \frac{r}{R} \right)^2 \right) \right]$$

# Formation of OHe-nucleus bound system



*Due to shielded Coulomb potential of X, Stark effect and tail of nuclear Yukawa force OHe attracts the nucleus. Nuclear attraction causes OHe excitation and Coulomb repulsion is switched on. If the system emits a photon, OHe forms a bound state with nucleus but **beyond** the nucleus with binding energy of few keV.*

# Few keV Level in OHe-nucleus system

- The problem is reduced to a quantum mechanical problem of energy level of OHe-nucleus bound state in the potential well, formed by shielded Coulomb, Stark effect and Yukawa tail attraction and dipole-like Coulomb barrier for the nucleus in vicinity of OHe. The internal well is determined by oscillatory potential of X in compound  $(Z+2)$  nucleus, in which He is aggregated.
- The numerical solution for this problem is simplified for rectangular wells and walls, giving a few keV level for Na.

# Rate of OHe-nucleus radiative capture

As soon as the energy of level is found one can use the analogy with radiative capture of neutron by proton with the account for:

- Absence of M1 transition for OHe-nucleus system (which is dominant for n+p reaction)
- Suppression of E1 transition by factor  $f \sim 10^{-3}$ , corresponding to isospin symmetry breaking

(in the case of OHe only isoscalar transition is possible, while E1 goes due to isovector transition only)

# Reproduction of DAMA/NaI and DAMA/LIBRA events

The rate of OHe radiative capture by nucleus with charge  $Z$  and atomic number  $A$  to the energy level  $E$  in the medium with temperature  $T$  is given by

$$\sigma v = \frac{f\pi\alpha}{m_p^2} \frac{3}{\sqrt{2}} \left(\frac{Z}{A}\right)^2 \frac{T}{\sqrt{Am_p E}}$$

Formation of OHe-nucleus bound system leads to energy release of its binding energy, detected as ionization signal. In the context of our approach the existence of annual modulations of this signal in the range 2-6 keV and absence of such effect at energies above 6 keV means that binding energy of Na-OHe system in DAMA experiment should not exceed 6 keV, being in the range 2-4 keV.



# Annual modulation of signals in DAMA/NaI and DAMA/LIBRA events

The amplitude of annual modulation of ionization signal (measured in counts per day per kg, cpd/kg) is given by

$$\zeta = \frac{3\pi\alpha \cdot n_0 N_A V_E t Q}{640\sqrt{2}A_{\text{med}}^{1/2}(A_I + A_{Na})} \frac{f}{S_3 m_p^2} \left(\frac{Z_i}{A_i}\right)^2 \frac{T}{\sqrt{A_i m_p E_i}} = 4.3 \cdot 10^{10} \frac{f}{S_3^2} \left(\frac{Z_i}{A_i}\right)^2 \frac{T}{\sqrt{A_i m_p E_i}}$$

This value should be compared with the integrated over energy bins signals in DAMA/NaI and DAMA/LIBRA experiments and the results of these experiments can be reproduced for

$$E_{Na} = 3keV$$

# **OPEN QUESTIONS OF THE OHE SCENARIO**

# Earth shadow effect

- OHe is nuclear interacting and thus should cause the Earth shadow effect.
- The studies, whether we can avoid recent DAMA constraints are under way.

# **THE PROBLEM OF POTENTIAL BARRIER**

# The crucial role of potential barrier in OHe-nucleus interaction

- Due to this barrier elastic OHe-nucleus scattering strongly dominates.
- If such barrier doesn't exist, overproduction of anomalous isotopes is inevitable.
- Its existence should be proved by proper quantum mechanical treatment

*J.-R. Cudell, M. Yu;Khlopov and Q.Wallemacq*

*Some Potential Problems of OHe Composite Dark Matter,*

*Bled Workshops in Physics (2014) V.15, PP.66-74; e-Print: arXiv: 1412.6030.*

**SENSITIVITY INDIRECT  
EFFECTS OF COMPOSITE  
DARK MATTER TO THE MASS  
OF THEIR DOUBLE CHARGED  
CONSTITUENTS**

# Excessive positrons in Integral

Taking into account that in the galactic bulge with radius  $\sim 1$  kpc the number density of O-helium can reach the value

$$n_o \approx 3 \cdot 10^{-3} / S_3 \text{ cm}^{-3}$$

one can estimate the collision rate of O-helium in this central region:

$$dN/dt = n_o^2 \sigma v_n 4\pi r_b^3 / 3 \approx 3 \cdot 10^{42} S_3^{-2} \text{ s}^{-1}$$

At the velocity of particules in halo, energy transfer in such collisions is  $E \sim 1$  MeV. These collisions can lead to excitation of O-helium. If 2S level is excited, pair production dominates over two-photon channel in the de-excitation by E0 transition and positron production with the rate

$$3 \cdot 10^{42} S_3^{-2} \text{ s}^{-1}$$

is not accompanied by strong gamma signal. This rate of positron production is sufficient to explain the excess of positron production in bulge, measured by Integral.

# Excessive positrons in Integral from dark atoms– high sensitivity to DM distribution




Figure 1: Values of the central dark matter density  $\rho_0$  ( $\text{GeV}/\text{cm}^3$ ) and of the OHe mass  $M$  (TeV) reproducing the excess of  $e^+e^-$  pairs production in the galactic bulge. Below the red curve, the predicted rate is too low.

*J.-R. Cudell, M. Yu. Khlopov and Q. Wallemacq*

*Dark atoms and the positron-annihilation-line excess in the galactic bulge.*

*Advances in High Energy Physics, vol. 2014, Article ID 869425, : arXiv: 1401.5228*



# Composite dark matter explanation for low energy positron excess

- In spite of large uncertainty of DM distribution in galactic bulge, where baryonic matter dominates and DM dynamical effects are suppressed, realistic simulations favor lower value of DM central density around  $\rho_0 \simeq 115 \text{ GeV/cm}^3$ . Then observed excess of positron annihilation line can be reproduced in OHe model only at the mass of its heavy double charged constituent:
  - $M \simeq 1.25 \text{ TeV}$

# A solution for cosmic positron excess?

- In WTC: if both technibaryons  $UU$  and technileptons  $\zeta$  are present, CDMS, LUX results constrain WIMP-like ( $UU\zeta$ ) component to contribute no more than 0,0001% of total DM density.
- Decays of positively charged  $UU \rightarrow l^+ l^+$  with a lifetime of about  $10^{21} s$  and mass 700-1000 GeV can explain the excess of cosmic positrons, observed by PAMELA and AMS02

# Cosmic positron excess from double charged constituents of dark atoms




Figure 3: Positron fraction in the cosmic rays from decays of dark matter particles (red curve), corresponding to the best-fit values of model parameters ( $M = 700$  GeV,  $\tau = 8 \cdot 10^{20}$  s,  $Br_{ee} = 0.182$ ,  $Br_{\mu\mu} = 0.394$ ,  $Br_{\tau\tau} = 0.424$ ), and fraction of secondary positrons (gray line), compared to the latest AMS-02 data [34] (blue dots).

*Probably such indirect effect is detected in the cosmic positron fluxes.*

*[figure from K.M.Belotsky et al. Int.J.Mod.Phys. D24 (2015) 1545004 arXiv:1508.02881 ]*

# Diffuse Gamma ray background




Figure 4: Gamma-ray flux multiplied by  $E^2$  from decays of dark matter particles in the Galaxy and beyond (green curve), corresponding to the best-fit values of model parameters ( $M = 700$  GeV,  $\tau = 8 \cdot 10^{20}$  s,  $Br_{ee} = 0.182$ ,  $Br_{\mu\mu} = 0.394$ ,  $Br_{\tau\tau} = 0.424$ ), compared to the latest FERMI/LAT data on isotropic diffuse gamma-ray background [42] ( $|b| > 20^\circ$ ,  $0^\circ \leq l < 360^\circ$  with point sources removed and without diffuse emission attributed to the interactions of Galactic cosmic rays with gas and radiation fields (foreground); here three different foreground models A (red dots), B (blue dots) and C (yellow dots) are shown). In our analysis we have used model B.


# Composite dark matter explanation for high energy positron excess

- Any source of high energy positrons, distributed in galactic halo is simultaneously the source of gamma ray background, measured by FERMI/LAT.
- Not to exceed the measured gamma ray background the mass of decaying double charged particles should not exceed

$$M < 1 \text{ TeV}$$

**COMPOSITE DARK MATTER  
CONSTITUENTS AT  
ACCELERATORS**

# Complementarity in searches for Dark Matter




*Usually, people use this illustration for complementarity in direct, indirect and accelerator searches for dark matter. However, we see that in the case of composite dark matter the situation is more nontrivial. We need charged particle searches to test dark atom model*

# Collider test for dark atoms

- Being the simplest dark atom model OHe scenario can not only explain the puzzles of direct dark matter searches, but also explain some possible observed indirect effects of dark matter. Such explanation implies a very narrow range of masses of (meta-) stable double charged particles in vicinity of 1 TeV, what is the challenge for their search at the experiments at the LHC.



# LHC discovery potential for charged components of composite dark matter



The shaded strips correspond to production cross sections of technileptons and A,C leptons with  $Q=2$  at  $7 \text{ teV} < \sqrt{s} < 14 \text{ TeV}$

# Search for multi-charge particles in the ATLAS experiment

Work is done in a frame of Multi-Charge Analysis Group

## Search for Multi-charge Objects in $pp$ collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector

K.M. Belotsky<sup>a</sup>, O. Bulekov<sup>a</sup>, M. Jünger<sup>b</sup>, M.Yu.Khlopov<sup>a,h</sup>, C. Marino<sup>c</sup>, P. Mermod<sup>d</sup>, H. Ogren<sup>e</sup>, A. Romaniouk<sup>a</sup>, Y. Smirnov<sup>a</sup>, W. Taylor<sup>f</sup>, B. Weinert<sup>g</sup>, D. Zieminska<sup>e</sup>, S. Zimmermann<sup>g</sup>

<sup>a</sup>*Moscow Engineering Physics Institute*

<sup>b</sup>*CERN*

<sup>c</sup>*University of Victoria*

<sup>d</sup>*Oxford University*

<sup>e</sup>*Indiana University*


<sup>f</sup>*York University*

<sup>g</sup>*University of Bonn*

<sup>h</sup>*University of Paris*

Our studies favor good chances for detection of multi-charge species in ATLAS detector

# Searches for multiple charged particles in ATLAS experiment



$M > 659 \text{ GeV}$   
 for  $|q|=2e$   
 at 95% c.l.  
 [Yu. Smirnov,  
 PhD Thesis]

[ATLAS Collaboration, Search for heavy long-lived multi-charged particles in pp collisions at  $\sqrt{s}=8 \text{ TeV}$  using the ATLAS detector, *Eur. Phys. J. C* 75 (2015) 362]

# Experimentum crucis for composite dark matter at the LHC

*Coming analysis of results of double charged particle searches at the LHC can cover all the range of masses, at which composite dark matter can explain excess of slow and high energy positrons, assuming the independent statistics in CMS and ATLAS experiments.*

Data period	Estimated lower mass limit, GeV	
	ATLAS or CMS separately	ATLAS and CMS combined
2015–2016	1000	1110
2015–2018	1190	1300

*Remind that composite dark matter can explain excess of low energy positrons at  $M=1.25$  TeV and high energy positrons at  $M<1$  TeV.*

# **PBH PROBES FOR PHYSICS OF EARLY UNIVERSE**

# Primordial Black Holes


- Any object of mass  $M$  can form Black hole, if contracted within its gravitational radius.

$$r \leq r_g = \frac{2GM}{c^2}$$

- It naturally happens in the result of evolution of massive stars (and, possibly, dense star clusters).
- In the early Universe Black hole can be formed, if expansion can stop within cosmological horizon [Zeldovich, Novikov, 1966]. It corresponds to strong nonhomogeneity in early Universe

$$\delta \equiv \frac{\delta\rho}{\rho} \sim 1$$

# Constraints on PBHs



# PBHs as indicator of early dust-like stages

- In homogeneous and isotropic Universe ( $\delta_0 \ll 1$ ) with equation of state  $p = k\varepsilon$  probability of strong nonhomogeneity  $\delta \sim 1$  is exponentially suppressed

$$P(\delta) = A(\delta, \delta_0) \exp\left(-\frac{k^2 \delta^2}{2\delta_0^2}\right)$$

- At  $k=0$  on dust-like stage exponential suppression is absent. The minimal estimation is determined by direct production of BHs


$$A(\delta, \delta_0) \geq \left(\frac{\delta_0}{\delta}\right)^5 \left(\frac{\delta_0}{\delta}\right)^{3/2} = \left(\frac{\delta_0}{\delta}\right)^{13/2}$$



# Dominance of superheavy particles

- Superheavy particles with mass  $m$  and relative concentration  $r = \frac{n}{n_\gamma}$  dominate in the Universe at  $T < r m$ .
- Coherent oscillations of massive scalar field also behave as medium with  $p=0$ .
- They form BHs either directly from collapse of symmetric and homogeneous configurations, or in the result of evolution of their gravitationally bound systems (pending on particle properties they are like « stars » or « galaxies »).


# PBHs as indicator of first order phase transitions



- Collision of bubbles with True Vacuum (TV) state during the first-order phase transition results in formation of False Vacuum (FV) bags, which contract and collapse in Black Holes (BH).

# PBH evaporation

- According to S. Hawking PBH with mass  $M$  evaporate due to creation of pairs by its nonstationary gravitational field. Products of evaporation have black body spectrum with



$$T_{PBH} \propto \frac{1}{r_g}$$

$$T_{PBH} \approx 10^{13} \text{ GeV} \left( \frac{1g}{M} \right)$$

- The rate of evaporation is given by

$$\frac{dM}{dt} = -\kappa T_{PBH}^4 r_g^2 \propto \frac{1}{r_g^2} \propto \frac{1}{M^2}$$

- The evaporation timescale is

$$t_{PBH} \approx 10^{27} \text{ s} \left( \frac{M}{1g} \right)^3$$

Any particle with  $m \leq T_{PBH}$

is created – UNIVERSAL source

# Effects of Primordial Black Holes

- PBHs behave like a specific form of Dark Matter
- Since in the early Universe the total mass within horizon is small, it seems natural to expect that such Primordial Black holes should have very small mass (much smaller, than the mass of stars). PBHs with mass  $M < 10^{15} g$  evaporate and their astrophysical effects are similar to effects of unstable particles.
- However, cosmological consequences of particle theory can lead to mechanisms of intermediate and even supermassive BH formation.

**STRONG PRIMORDIAL  
INHOMOGENEITY PROBES  
FOR INFLATION AND  
BARYOSYNTHESIS**

# Strong nonhomogeneities in nearly homogeneous and isotropic Universe

- The standard approach is to consider homogeneous and isotropic world and to explain development of nonhomogeneous structures by gravitational instability, arising from small initial fluctuations.

$$\delta \equiv \delta\rho / \rho \ll 1$$

- However, if there is a tiny component, giving small contribution to total  $\rho_i \ll \rho$  its strong nonhomogeneity  $\delta_i \equiv (\delta\rho / \rho)_i > 1$

is compatible with small nonhomogeneity of the total density

$$\delta = (\delta\rho_i + \delta\rho) / \rho \approx (\delta\rho_i / \rho_i)(\rho_i / \rho) \ll 1$$

Such components naturally arise as consequences of particle theory, shedding new light on galaxy formation and reflecting in cosmic structures the fundamental structure of microworld.

# Strong Primordial nonhomogeneities from the early Universe

- Cosmological **phase transitions** in inflationary Universe can give rise to unstable cosmological defects, retaining a replica in the form of primordial **nonlinear** structures (massive PBH clusters, archioles).
- Nonhomogenous baryosynthesis (including spontaneous baryosynthesis and leptogenesis) in its extreme form can lead to **antimatter** domains in baryon asymmetrical inflationary Universe.

Strong nonhomogeneities of total density and baryon density are severely constrained by CMB data at large scales (and by the observed gamma ray background in the case of antimatter). However, their existence at smaller scales is possible.

# Cosmological Phase transitions 1.

- At high temperature  $T > T_{cr}$  spontaneously broken symmetry is restored, owing to thermal corrections to Higgs potential

$$V(\varphi, T = 0) = -\frac{m^2}{2}\varphi^2 + \frac{\lambda}{4}\varphi^4 \Rightarrow V(\varphi, T) = \left( C\lambda T^2 - \frac{m^2}{2} \right) \varphi^2 + \frac{\lambda}{4}\varphi^4$$

- When temperature falls down below

$$T = T_{cr} \cong \langle \varphi \rangle = \frac{m}{\sqrt{\lambda}}$$

transition to phase with broken symmetry takes place.



# Cosmological Phase transitions 2.

- Spontaneously broken symmetry can be restored on chaotic inflationary stage, owing to corrections in Higgs potential due to interaction of Higgs field with inflaton

$$V(\varphi, \psi = 0) = -\frac{m^2}{2} \varphi^2 + \frac{\lambda}{4} \varphi^4 \Rightarrow V(\varphi, \psi) = \left( \varepsilon \psi^2 - \frac{m^2}{2} \right) \varphi^2 + \frac{\lambda}{4} \varphi^4$$

- When inflaton field rolls down below

$$\psi = \psi_{cr} \cong \frac{m}{\sqrt{\varepsilon}}$$

transition to phase with broken symmetry takes place.

# Topological defects

- In cosmological phase transition false (symmetric) vacuum goes to true vacuum with broken symmetry. Degeneracy of true vacuum states results in formation of topological defects.
- Discrete symmetry of true vacuum  $\langle \varphi \rangle = \pm f$  leads to domains of true vacuum with  $+f$  and  $-f$  and false vacuum wall on the border.
- Continuous degeneracy  $\langle \varphi \rangle = f \exp(i\theta)$  results in succession of singular points surrounded by closed paths with  $\Delta\theta = 2\pi$ . Geometrical place of these points is line – cosmic string.
- SU(2) degeneracy results in isolated singular points – in GUTs they have properties of magnetic monopoles.

# U(1) model

$$V(\psi) = \frac{\lambda}{2} (\psi^2 - f^2)^2$$

**After spontaneous symmetry breaking infinitely degenerated vacuum**

$$\psi = f e^{i\varphi/f}$$

**experiences second phase transition due to the presence  
(or generation by instanton effects)**


$$V(\varphi) = \Lambda^4 (1 - \cos(\varphi/f))$$

**to vacuum states**

$$\theta \equiv \varphi/f = 0, 2\pi, \dots$$

**In particular, this succession of phase transitions takes place in axion models**

# Topological defects




- Spontaneous breaking of U(1) symmetry results in the continuous degeneracy of vacua. In the early Universe the transition to phase with broken symmetry leads to formation of cosmic string network.
- The tilt in potential breaks continuous degeneracy of vacua. In the result string network converts into walls-bounded-by-strings structure in the second phase transition. This structure is unstable and decay, but the initial values of phase define the energy density of field oscillations.

# Unstable topological defects

- This picture takes place in axion cosmology.
- The first phase transition gives rise to cosmic axion string network.
- This network converts in the second phase transition into walls-bounded-by-strings structure (walls are formed between strings along the surfaces  $\alpha = \pi$ ), which is unstable.
- However, the energy density distribution of coherent oscillations of the field  $\alpha$  follows the walls-bounded-by-strings structure.

# Archioles structure




- Numerical studies revealed that ~80% of string length corresponds to infinite Brownian lines, while the remaining ~20% of this length corresponds to closed loops with large size loops being strongly suppressed. It corresponds to the well known scale free distribution of cosmic strings.
- The fact that the energy density of coherent axion field oscillations reflects this property is much less known. It leads to a large scale correlation in this distribution, called archioles.
- Archioles offer possible seeds for large scale structure formation.
- However, the observed level of isotropy of CMB puts constraints on contribution of archioles to the total density and thus puts severe constraints on axions as dominant form of Dark Matter.

# Massive Primordial Black Holes

- Any object can form Black hole, if contracted within its gravitational radius. It naturally happens in the result of evolution of massive stars (and, possibly, star clusters).
- In the early Universe Black hole can be formed, if within cosmological horizon expansion can stop [Zeldovich, Novikov, 1966]. Since in the early Universe the total mass within horizon is small, it seems natural to expect that such Primordial Black holes should have very small mass (much smaller, than the mass of stars).
- However, cosmological consequences of particle theory can lead to mechanisms of intermediate and even supermassive BH formation.

# Closed walls formation in Inflationary Universe



If the first U(1) phase transition takes place on inflationary stage, the value of phase  $\theta$ , corresponding to e-folding  $N \sim 60$ , fluctuates



$$\Delta\theta \approx H_{\text{infl}} / (2\pi f)$$

Such fluctuations can cross  $\pi$

and after coherent oscillations begin, regions with  $\theta > \pi$  occupying relatively small fraction of total volume are surrounded by massive walls



# Massive PBH clusters



Each massive closed wall is accompanied by a set of smaller walls.

As soon as wall enters horizon, it contracts and collapses in BH.

Each locally most massive BH is accompanied by a cloud of less massive BHs.

The structure of such massive PBH clouds can play the role of seeds for galaxies and their large scale distribution.

# Spectrum of Massive BHs

- The minimal mass of BHs is given by the condition that its gravitational radius exceeds the width of wall ( $d \approx 2f/\Lambda^2$ )

$$r_g = \frac{2M}{m_{Pl}^2} > d = \frac{2f}{\Lambda^2} \Rightarrow M_{\min} = f \left( \frac{m_{Pl}}{\Lambda} \right)^2$$

- The maximal mass is given by the condition that pieces of wall do not dominate within horizon, before the whole wall enters the horizon

$$R < \frac{3\sigma_w}{\rho_{tot}} \Rightarrow M_{\max} = f \left( \frac{m_{Pl}}{f} \right)^2 \left( \frac{m_{Pl}}{\Lambda} \right)^2 \Rightarrow \frac{M_{\max}}{M_{\min}} = \left( \frac{m_{Pl}}{f} \right)^2$$

# GW signals from closed wall collapse and BHs merging in clouds

- Closed wall collapse leads to primordial GW spectrum, peaked at  $\nu_0 = 3 \cdot 10^{11} (\Lambda/f) \text{ Hz}$  with energy density up to



$$\Omega_{GW} \approx 10^{-4} (f/m_{Pl})$$

- At  $f \sim 10^{14} \text{ GeV}$   $\Omega_{GW} \sim 10^{-9}$
- For  $1 < \Lambda < 10^8 \text{ GeV}$   $3 \cdot 10^{-3} \text{ Hz} < \nu_0 < 3 \cdot 10^5 \text{ Hz}$
- Merging of BHs in BH cluster is probably detected by LIGO!.

# The first GW signal!


14th September 2015 - GW150914

2



From VIA talk 09.12.2016 by P.Lasky

# Abbott et al. (2016) - GW150914




# GW astronomy!

## 2.5 measurements!


9

September 2015 — February 2016



Abbott et al. (2016) - O1 BBH

# GW Astronomy: Signal 170814 was registered by 3 detectors!



*Joint detection of the GW signal 170814 by two LIGO detectors and VIRGO detector can provide localization of the source!*

# Binaries of massive PBHs?

- Massive PBHs are not distributed homogeneously in space, but are in clouds.
- It makes more probable formation of massive PBHs binaries.
- The problem of creation of stellar mass PBH clouds, their evolution and formation of BH binaries in them may be an interesting hot topic for a PhD thesis



**ANTIMATTER STARS IN  
BARYON ASYMMETRIC  
UNIVERSE**

# Antimatter from nonhomogeneous baryosynthesis

- Baryon excess  $B > 0$  can be generated nonhomogeneously  $B(x)$ .
- Any nonhomogeneous mechanism of BARYON excess generation  $B(x)$  leads in extreme form to ANTIBARYON excess in some regions.

# Survival of antimatter domains

Diffusion of baryons and antibaryons to the border of domain results in eating of antimatter by surrounding baryonic matter.



$$\partial n_b / \partial t = D(t) \partial^2 n_b / \partial x^2 - \alpha n_b \quad \text{where} \quad D(t) \approx \frac{3T_p c}{2\rho_\gamma \sigma_T}$$

The minimal surviving scale is given by

$$d \approx \frac{c}{\sqrt{\frac{8\pi}{3} G \rho_0}} \frac{T_p}{m} \sqrt{\frac{m}{T_{rec}}} \int_{T_p/T_{rec}}^1 \frac{dy}{y^{3/2}} = \frac{2c}{\sqrt{\frac{8\pi}{3} G \rho_0}} \sqrt{\frac{T_p}{m}}$$

which is about  $d \sim 3/h$  kpc.

# Nonhomogeneous spontaneous baryosynthesis



- Model of spontaneous baryosynthesis provides quantitative description of combined effects of inflation and nonhomogeneous baryosynthesis, leading to formation of antimatter domains, surviving to the present time.



# Antimatter in galaxies

Number of e-fold	Number of domains	Size of domain
59	0	1103Mpc
55	$5.005 \cdot 10^{-14}$	37.7Mpc
54	$7.91 \cdot 10^{-10}$	13.9Mpc
52	$1.291 \cdot 10^{-3}$	1.9Mpc
51	0.499	630kpc
50	74.099	255kpc
49	$8.966 \cdot 10^3$	94kpc
48	$8.012 \cdot 10^3$	35kpc
47	$5.672 \cdot 10^7$	12kpc
46	$3.345 \cdot 10^9$	4.7kpc
45	$1.705 \cdot 10^{11}$	1.7kpc

**Numerical simulations show that within the modern horizon possible amount of antimatter domains, with the size exceeding the survival scale and thus surviving to the present time, can be comparable with the total number of galaxies.**


**In our Galaxy from 1000 to 100000 antimatter stars can exist in a form of antimatter globular cluster (Khlopov, 1998). Being in halo, such cluster is a faint gamma ray source, but antimatter from it pollutes Galaxy and can be observed indirectly by annihilation, or directly as anti-meteorites or antinuclei in cosmic rays.**

# Antimatter pollution of Galaxy




- Since antihydrogen is dominant in antimatter composition, the Galaxy is dominantly polluted by antiprotons.
- Their lifetime in Galaxy depends on their velocity and density of surrounding matter.

# Gamma background from antimatter annihilation in Galaxy



- Antiproton annihilation can reproduce gamma background observed by EGRET in the range tens-hundreds MeV.
- It can not be considered as PROOF for existence of antimatter stars – only pieces of antimatter (antihelium nuclei, antimeteorites) can provide such PROOF.

# Cosmic antihelium test for antimatter stars in Galaxy



- **Nonhomogeneous baryosynthesis in extreme form leads to antimatter domains in baryon asymmetrical Universe**
- **To survive in the surrounding matter domain should be sufficiently large, and to have sufficiently high internal antibaryon density to form stars. It gives minimal estimation of possible amount of antimatter stars in Galaxy**
- **The upper limit comes from observed gamma background**
- **Assuming that antihelium component of cosmic rays is proportional to the fraction of antimatter stars in the total mass of Galaxy, it is possible to test this hypothesis initially in PAMELA and then completely in AMS-02 experiment**



# First signal from antimatter stars in AMS02?




Figure 14. An antihelium candidate.

*Presented in CERN on 08.12.2016 by Prof. S.Ting*

# Latest Results from the AMS Experiment on the International Space Station

## ABSTRACT


In seven years on the Space Station, AMS has collected more than 115 billion charged cosmic rays with energies up to multi TeV. The measured positron spectra agrees well with dark matter models. The energy dependence of elementary particles (electrons, positrons, protons and antiprotons) as well as the rigidity dependence of primary cosmic rays and secondary cosmic rays are unique and distinct. These results require a new understanding of the cosmos.

*Samuel Ting*  
*24 May 2018*

# Antihelium events

- 8 clear single track events with  $Z=-2$  within helium mass region
- Momentum resolution better than 10%

# Antihelium-4 candidates!



*Samuel Ting 24 May 2018*

## Puzzle of antiHe-3 and antiHe-4 ratio

Two anti-Helium-4 events are announced on 24.05.2018 with background probability 1/300.

Continuing to take the data through 2024 would reduce background probability, putting such candidate events above 5-sigma significance

Though He3/He4 ratio is 0.1-0.2, the antiHe3/antiHe4 ratio looks now like 3. More data will resolve this puzzle

# Conclusions

- **Physical basis of the modern cosmology implies new physics.**
- **Dark atom hypothesis can explain puzzles of direct dark matter searches and cosmic positron anomalies. It can be directly probed at the LHC.**
- **PBHs provide probes for physics of very early Universe**
- **Strong primordial nonlinear structures (massive PBH clouds, strong nonhomogeneities of baryonic matter and even antimatter stars) provide cosmological probes for physics of inflationary models with baryosynthesis and dark matter.**