

How far has so far the Spin-Charge-Family theory succeeded to offer the explanation for the observed phenomena :

- i. The standard model assumptions,**
- ii. second quantized fermion fields,**
- iii. matter-antimatter asymmetry,**
- iv. the appearance of the dark matter,.....,**

making several predictions

N.S. Mankoč Borštnik, University of Ljubljana
24rd International Workshop "What comes beyond the standard models?"

Some publications:

- ▶ *Phys. Lett.* **B 292**, 25-29 (1992), *J. Math. Phys.* **34**, 3731-3745 (1993), *Mod. Phys. Lett.* **A 10**, 587-595 (1995), *Int. J. Theor. Phys.* **40**, 315-337 (2001),
- ▶ *Phys. Rev.* **D 62** (04010-14) (2000), *Phys. Lett.* **B 633** (2006) 771-775, **B 644** (2007) 198-202, **B** (2008) 110.1016, *JHEP* **04** (2014) 165, *Fortschritte Der Physik-Progress in Physics*, (2017) with H.B.Nielsen,
- ▶ *Phys. Rev.* **D 74** 073013-16 (2006), with A.Borštnik Bračič,
- ▶ *New J. of Phys.* **10** (2008) 093002, arxiv:1412.5866, with G.Bregar, M.Breskvar, D.Lukman,
- ▶ *Phys. Rev.* **D** (2009) 80.083534, with G. Bregar,
- ▶ *New J. of Phys.* (2011) 103027, *J. Phys. A: Math. Theor.* **45** (2012) 465401, *J. Phys. A: Math. Theor.* **45** (2012) 465401, *J. of Mod. Phys.* **4** (2013) 823-847, arxiv:1409.4981, **6** (2015) 2244-2247, *Phys. Rev.* **D 91** (2015) 6, 065004, . *J. Phys.: Conf. Ser.* **845 01 IARD 2017**, *Eur. Phys. J.C.* **77** (2017) 231, Rev. Article to appear in *Prog. in Part. and Nucl. Phys.* **PPNP 2021**,

More than **50 years ago** the **electroweak (and colour) standard model** offered an **elegant new step** in **understanding the origin of fermions and bosons** by **postulating**:

A.

- ▶ The existence of **massless family members** with the **charges** in the **fundamental representation of the groups** -
 - the **coloured triplet quarks and colourless leptons**,
 - the **left handed members as the weak charged doublets**,
 - the **right handed weak chargeless members**,
 - the **left handed quarks distinguishing in the hyper charge from the left handed leptons**,
 - **each right handed member having a different hyper charge**.
- ▶ The existence of **massless families to each of a family member**.

α name	hand- edness $-4iS^0 S^{12}$	weak charge τ^{13}	hyper charge Y	colour charge	elm charge Q
u_L^i	-1	$\frac{1}{2}$	$\frac{1}{6}$	colour triplet	$\frac{2}{3}$
d_L^i	-1	$-\frac{1}{2}$	$\frac{1}{6}$	colour triplet	$-\frac{1}{3}$
ν_L^i	-1	$\frac{1}{2}$	$-\frac{1}{2}$	colourless	0
e_L^i	-1	$-\frac{1}{2}$	$-\frac{1}{2}$	colourless	-1
u_R^i	1	weakless	$\frac{2}{3}$	colour triplet	$\frac{2}{3}$
d_R^i	1	weakless	$-\frac{1}{3}$	colour triplet	$-\frac{1}{3}$
ν_R^i	1	weakless	0	colourless	0
e_R^i	1	weakless	-1	colourless	-1

Members of each of the $i = 1, 2, 3$ families, $i = 1, 2, 3$ massless before the electroweak break. Each family contains the left handed weak charged quarks and the right handed weak chargeless quarks, belonging to the colour triplet $(1/2, 1/(2\sqrt{3}))$, $(-1/2, 1/(2\sqrt{3}))$, $(0, -1/(\sqrt{3}))$.

And the anti-fermions to each family and family member.

B.

- ▶ The existence of **massless vector gauge fields** to the observed **charges** of the **family members**, **carrying charges** in the **adjoint representation of the charge groups**.

Masslessness needed for gauge invariance.

Gauge fields before the electroweak break

- ▶ Three massless vector fields, the gauge fields of the three charges.

name	hand- edness	weak charge	hyper charge	colour charge	elm charge
hyper photon	0	0	0	colourless	0
weak bosons	0	triplet	0	colourless	triplet
gluons	0	0	0	colour octet	0

They all are vectors in $d = (3 + 1)$, in the adjoint representations with respect to the weak, colour and hyper charges.

C.

- ▶ The **existence of a massive scalar field - the higgs**,
 - carrying the weak charge $\pm\frac{1}{2}$ and the hyper charge $\mp\frac{1}{2}$ (as it would be in the fundamental representation of the groups.)
 - gaining at some step the **imaginary mass** and consequently the **constant value** , breaking the weak and the hyper charge and correspondingly breaking the **mass protection**.
- ▶ The **existence** of the **Yukawa couplings**, taking care of
 - the properties of **fermions** and
 - the masses of the **heavy bosons**.

- ▶ The **Higgs's field**, the scalar in $d = (3 + 1)$, a **doublet with respect to the weak charge**.

name	hand- edness	weak charge	hyper charge	colour charge	elm charge
0. $Higgs_u$	0	$\frac{1}{2}$	$\frac{1}{2}$	colourless	1
$\langle Higgs_d \rangle$	0	$-\frac{1}{2}$	$\frac{1}{2}$	colourless	0

name	hand- edness	weak charge	hyper charge	colour charge	elm charge
$\langle Higgs_u \rangle$	0	$\frac{1}{2}$	$-\frac{1}{2}$	colourless	0
0. $Higgs_d$	0	$-\frac{1}{2}$	$-\frac{1}{2}$	colourless	-1

D.

- ▶ There is the **gravitational field** in **$d=(3+1)$** .

- ▶ **The *standard model* assumptions have been confirmed without offering surprises.**
- ▶ The last unobserved field as a field, the **Higgs's scalar**, detected in June 2012, was confirmed in March 2013.
- ▶ The waves of the **gravitational field** were detected in February 2016 and again 2017.

There remain not understood phenomena:

- ▶ The *Standard model* assumptions **need explanation**.
- ▶ **There are several cosmological observations which do not look to be explainable within the *standard model*, like**
 - The **existence of the *dark matter***
 - The ***matter/antimatter asymmetry* in the universe**
 - The **need for the *dark energy***
- ▶ the **observed dimension of space time**,
- ▶ the **quantization of the gravitational field**,
- ▶ ...

- ▶ The *standard model* assumptions have in the literature several explanations, but with **many new not explained assumptions**.

- ▶ I am proposing **the *spin-charge-family* theory**, which offers the explanation for
 - i. all the assumptions of the *standard model*,
 - ii. for many observed phenomena:
 - ii.a. the **dark matter**,
 - ii.b. the **matter-antimatter** asymmetry,
 - ii.c. **others observed phenomena**,
 - iii. explaining the Dirac's postulates for the second quantized fermions,
 - iv. **making several predictions**.

Is this the right next step beyond both *standard models*?

- ▶ Work done so far on the **spin-charge-family theory** is promising.

**** We try to understand:**

- ▶ **What are elementary constituents and interactions among constituents in our Universe, in any universe?**
- ▶ **Can the elementary constituent be of only one kind? Are the four observed interactions — gravitational, elektromagnetic, weak and colour — of the common origin?**
- ▶ **Is the space-time the so far observed (3 + 1)? Why (3+1)?**
- ▶ **If not (3 + 1) may it be that the space-time is infinite?**
- ▶ **How has the space-time of our universe started?**
- ▶ **What is the matter and what the anti-matter?**

Obviously it is the time to make **the next step beyond both standard models.**

What questions should one ask to be able to find **next steps** beyond the *standard models* and to understand not yet understood phenomena?

- ▶ ○ Where do **family members** originate?
 - Where do **charges** of **family members** originate?
 - Why are the **charges** of **family members** so different?
 - Why have the **left handed family members** so different charges from the **right handed** ones?
- ▶ ○ Where do **families** of **family members** originate?
 - How **many different families** exist?
 - Why do **family members – quarks and leptons** – manifest so different properties if they all start as massless?

- ▶ **o** How is the **origin** of the **scalar field** (the Higgs's scalar) and the **Yukawa couplings connected** with the origin of **families**?
 - o** How many **scalar fields** determine properties of the so far (and others possibly be) **observed fermions** and masses of **weak bosons**? (The Yukawa couplings certainly speak for the existence of several scalar fields with the properties of Higgs's scalar.)
- ▶ **Why is the Higgs's scalar**, or are all **scalar fields**, if there are several, **doublets** with respect to the weak and the hyper charge?
- ▶ **Do exist** also **scalar fields** with the **colour charge in the fundamental representation** and where, if they are, **do they manifest**?

- ▶ Where do the **charges** and correspondingly the so far (and others possibly be) **observed vector gauge fields** originate?
- ▶ **Where** does the **dark matter** originate?
- ▶ **Where** does the "ordinary" **matter-antimatter asymmetry originate?**
- ▶ **Where** does the **dark energy** originate?
- ▶ What is the dimension of space? $(3 + 1)?$, $((d - 1) + 1)?$, $\infty?$
- ▶ **What** is the role of the **symmetries**– discrete, continuous, global and gauge – in our **universe, in Nature?**
- ▶ And many others.

My statement:

- ▶ **An elegant trustworthy next step** must offer answers to **several** open questions, explaining:
 - o The **origin of the family members and the charges.**
 - o The **origin of the families and their properties.**
 - o The **origin of the scalar fields and their properties.**
 - o The **origin of the vector fields and their properties.**
 - o The **origin of the dark matter.**
 - o The **origin of the "ordinary" matter-antimatter asymmetry.**

My statement continues:

- ▶ There exist not yet observed families, gauge vector and scalar gauge fields.
- ▶ **Dimension of space is larger than 4** (very probably infinite).
- ▶ Inventing a next step which covers one of the open questions, might be of a help **but can hardly show the right next step in understanding nature.**

In the literature **NO explanation for the existence of the families can be found**, which would not just assume the family groups.

Several extensions of the **standard model** are, however, proposed, like:

- ▶ The $SU(3)$ group is assumed to describe – not explain – the existence of three families.

Like the **Higgs's** scalar charges are in the **fundamental** representations of the groups, also the **Yukawas** are assumed to emerge from the scalar fields, in the **fundamental** representation of the $SU(3)$ group.

- ▶ **SU(5) and SO(10) grand unified theories are proposed, unifying all the charges.** But the **spin** (the handedness) is obviously connected with the (weak and the hyper) charges, what these theories do "by hand" as it does the *standard model*, and the appearance of families is not explained.
- ▶ **Supersymmetric theories, assuming the existence of bosons with the charges of quarks and leptons and fermions with the charges of the gauge vector fields, although having several nice properties but not explaining the appearance of families (except again by assuming larger groups), are not, to my understanding, the right next step beyond the *standard model*.**

- o The **Spin-Charge-Family** theory does offer the **explanation for all the assumptions of the standard model**, answering up to now several of the above cited open questions!
 - o The **more effort** is put into this theory, the more answers to the open questions in elementary particle physics and cosmology is the theory offering.

- o I shall first make a short introduction into the **Spin-Charge-Family** theory.
- o I shall make an overview of achievements so far of the **Spin-Charge-Family** theory.
- o I shall report on **how does the odd Clifford algebra explain the second quantization postulates of Dirac.**
[arXiv:1802.05554v1v2, arXiv:1802.05554v4, arXiv:1902.10628]
Rev. article to appear in **JPPNP –2021** Progress in Particle and Nuclear Physics.

- ▶ A brief introduction into the **spin-charge-family theory**.

- ▶ **Spinors** carry in $d \geq (13 + 1)$ **two kinds of spin, no charges**, *Phys. Rev. D* **91** 065004 (2015), *J.of Mod. Physics* **6** (2015) 2244, **Rev. article** in **JPPNP** to appear in 2021.
 - The **Dirac spin** (γ^a) in $d = (13 + 1)$ **describes** in $d = (3 + 1)$ **spin** and **ALL the charges of quarks and leptons, left and right handed**, explaining all the **assumptions about the charges and the handedness of the standard model** *J. of Math. Phys.* **34** (1993), 3731, *J. of Math. Phys.* **43**, 5782 (2002) [hep-th/0111257].
 - The **second kind of spin** ($\tilde{\gamma}^a$) **describes FAMILIES**, explaining the **origin and numbers of families**, *J. of Math. Phys.* **44** 4817 (2003) [hep-th/0303224].
 - There is **NO third kind of spin**.
- ▶ **C,P,T symmetries** in $d = (3 + 1)$ **follow from the C,P,T symmetry** in $d \geq (13 + 1)$. (*JHEP* **04** (2014) 165)

- ▶ All **vector** and **scalar gauge fields** origin in gravity, explaining the origin of the vector and scalar gauge fields, which in the *standard model* are assumed *Eur. Phys. J. C 77* (2017) 231:
 - o **Vector** and **scalar gauge fields** origin in two **spin connection fields**, the **gauge fields** of $\gamma^a\gamma^b$ and $\tilde{\gamma}^a\tilde{\gamma}^b$, and in
 - o **vielbeins**, the gauge fields of momenta
Eur. Phys. J. C 77 (2017) 231, [arXiv:1604.00675]

- ▶ If there are **no spinor sources present**, then either vector (\vec{A}_m^A , $m = 0, 1, 2, 3$) or scalar (\vec{A}_s^A , $s = 5, 6, \dots, d$) gauge fields are determined by **vielbeins** uniquely.

- ▶ **Spinors** interact correspondingly with
 - the **vielbeins** and
 - the **two kinds of the spin connection fields**, *Eur. Phys. J. C* **77** (2017) 231.
- ▶ In $d = (3 + 1)$ the **spin-connection fields**, together with the **vielbeins**, manifest either as
 - **vector** gauge fields with all the **charges** in the **adjoint** representations or as
 - **scalar** gauge fields with the **charges** with respect to the **space index** in the **"fundamental"** representations and all the other **charges** in the **adjoint** representations or as
 - **tensor** gravitational field.

There are two kinds of **scalar fields** with respect to the space index s :

- ▶ Those with ($s = 5, 6, 7, 8$) (they carry zero "**spinor charge**") are **doublets** with respect to the $SU(2)_I$ (the weak) charge and the **second $SU(2)_{II}$ charge** (determining the hyper charge). They are in the **adjoint** representations with respect to the **family** and the **family members charges**.
- These **scalars** explain the **Higgs's scalar** and the **Yukawa couplings**.

Phys. Rev. **D 91** (2015) 6, 065004

- ▶ **Those** with twice the "spinor charge" of a quark and ($s = 9, 10, ..d$) are **colour triplets**. Also they are in the **adjoint representations** with respect to the **family** and the **family members charges**.
 - These **scalars** transform **antileptons** into **quarks**, and **antiquarks** into **quarks** and back and correspondingly **contribute to matter-antimatter asymmetry** of our universe and to **proton decay**.
- ▶ There are **no additional scalar fields** in the **spin-charge-family theory**, if $d = (13 + 1)$.

Phys. Rev. **D 91** (2015) 6, 065004

J. of Mod. Phys. **6** (2015) 2244

Breaking symmetry from M^{13+1} into $M^{7+1} \times M^6$

- ▶ We start with the massless solutions of the Weyl equation in $d = (13 + 1)$ with the "basis vectors", described by the odd Clifford algebra objects, determining the internal space of fermions.
- ▶ With the spin (or the total angular momentum) in extra dimensions, $d > (7 + 1)$, determining the charge in $d = (7 + 1)$.
- ▶ We then let the \mathcal{M}^{13+1} manifold to break into $\mathcal{M}^{7+1} \times$ an almost S^6 sphere.
- ▶ The Weyl equation, $m = (0, 1, 2, 3, 5, 6, 7, 8)$ and $s = 9, 10, \dots, 13, 14$ is

$$(\gamma^m p_m + \gamma^s p_{0s})\psi = 0,$$
$$p_{0s} = f_s^\sigma (p_\sigma - \frac{1}{2} S^{ab} \omega_{ab\sigma} - \frac{1}{2} \tilde{S}^{ab} \tilde{\omega}_{ab\sigma}) + \frac{1}{2E} \{p_\sigma, f_s^\sigma E\}_-.$$

- ▶ With the choice of the vielbein fields and the spin spinconnection fields of both kinds one can achieve that the infinite surface $d = (9, 10, 11, \dots, 13, 14)$ curls into an almost S^6 (with one hole with the substructure of $SU(3) \times U(1)$) with massless fermions in $d = (7 + 1)$.
- ▶ This is the project, not yet done. The simpler problem with breaking \mathcal{M}^{5+1} manifold into $\mathcal{M}^{3+1} \times$ an almost S^2 sphere with one hole is done, without taking into account families and with families included.

New J. Phys. 13:103027, 2011.

J. Phys. A: Math. Theor. 45:465401, 2012.

Condensate

- ▶ The (assumed so far, waiting to be derived how does this spontaneously appear) **scalar condensate** of **two right handed neutrinos** with the **family** quantum numbers of the upper four families (there are two four family groups in the theory), appearing $\approx 10^{16}$ GeV or higher,
 - **breaks the CP** symmetry, causing the **matter-antimatter asymmetry** and the proton decay,
 - couples to all the **scalar fields**, making them massive,
 - couples to all the phenomenologically **unobserved vector gauge fields**, making them massive.
 - Before the electroweak break all the so far **observed vector gauge fields are massless**.

Phys. Rev. **D 91** (2015) 6, 065004,

J. of Mod. Phys. **6** (2015) 2244,

J. Phys.: Conf.Ser. 845 01, **IARD 2017**

- ▶ The **vector fields**, which do not couple to the condensate and remain massless, are:
 - o the **hyper charge vector field**.
 - o the **weak vector fields**,
 - o the **colour vector fields**,
 - o the **gravity fields**.

The $SU(2)_{II}$ symmetry breaks due to the **condensate**, leaving the **hyper charge unbroken**.

Nonzero vacuum expectation values of scalars

— waiting to be shown how does such an event, making the masses of the scalar gauge fields imaginary, appear in the *spin-charge-family* spontaneously.

- ▶ The scalar fields with the **space index (7, 8)**, gaining **nonzero vacuum expectation values**, a constant values, cause the **electroweak break**,
 - breaking the weak and the hyper charge,
 - changing their own masses,
 - bringing masses to the **weak bosons**,
 - bringing masses to the **families of quarks and leptons**.

Phys. Rev. **D 91** (2015) 6, 065004,

J. Phys.: Conf.Ser. 845 01 **IARD 2017**,

Eur. Phys. J.C. **77** (2017) 231 [arXiv:1604.00675],

J. of Mod. Phys. **6** (2015) 2244, [arXiv:1502.06786,

arXiv:1409.4981]

- ▶ The only gauge fields which do not couple to these scalars and remain massless are
 - electromagnetic,
 - colour vector gauge fields,
 - gravity.
- ▶ There are two times four decoupled massive **families** of **quarks and leptons** after the electroweak break:
 - There are the observed **three families** among the lower **four, the fourth to be observed**.
 - The stable among the **upper four families** form the **dark matter**.

Phys. Rev. **D 80**, 083534 (2009),

Phys. Rev. **D 91** (2015) 6, 065004,

J. Phys.: Conf.Ser. 845 01, **IARD 2017**

- ▶ It is **extremely encouraging** for the **spin-charge-family theory**, that a **simple starting action** contains **all the degrees of freedom observed at low energies**, directly or indirectly, and that only
 - the **break of manifold** $M^{(13,1)}$ to $M^{(7,1)} \times M^{(6)}$ is needed so that the manifold $M^{(6)}$ makes an almost S^n sphere.
 - the **condensate** and
 - **constant values of all the scalar fields with $s = (7, 8)$** are needed that the **theory explains**
 - **all the assumptions** of the standard model, with the gauge fields, scalar fields, families of fermions, masses of fermions and of bosons included,
 - explaining also **the dark matter**,
 - **the matter/antimatter asymmetry**,
 - **the triangle anomalies cancellation** in the standard model (Forts. der Physik, Prog.of Phys.) (2017) 1700046) and...

The spin-charge-family theory

The **spin-charge-family** theory is a kind of a **Kaluza-Klein-like** theory, but with **two kinds of spins**.

In d -dimensional space there are **fermions** with **two kinds of spins** and **gravity**, represented by two **spin connection** and **vielbein gauge fields**.

J. of Mod. Phys. **4** (2013) 823,
Phys. Rev. **D 91** (2015) 6, 065004,
J. of Mod. Phys. **6** (2015) 2244 [arxiv:1409.4981],
IARD, J. Phys.: Conf. Ser. 845 (2017) 012017
[arXiv:1607.01618/v2],
Eur. Phys. J.C. **77** (2017) 231,
Forts. der Physik, Prog. of Phys.) (2017) 1700046
Rev. art. in Prog. Part. Nucl. Phys. 2021.

Comparison of the **spin-charge-family** theory and the **unifying theories**, with the **Kaluza-Klein like** theories included.

- ▶ The $SO(10)$ group is the subgroup of $SO(13, 1)$ group, but only if taking into account the whole group the handedness in $d = (3 + 1)$ is related to charges, otherwise this relation must be made by "hand".
- ▶ The $SO(10)$ -unifying theories assume the existence of all the corresponding gauge fields. The family group must be assumed in addition.
- ▶ The **spin-charge-family** theory has in the simple starting action **only gravity** as **gauge fields**.

- ▶ The recognition that there are **two kinds of the Clifford algebras** objects,
 - one **used to describe spins and charges** of fermions,
 - the second one **describing families** of fermions,
 - the theory **unifies spins, charges and families**.
- ▶ Paying attention that the **odd Clifford algebra** , describing internal space of fermions, makes creation and annihilation operators **anticommuting objects**, the theory offers the **second quantized fermion fields without Dirac's postulates**.
- ▶ **At low energies the simple starting action** for fermions and **gravity** manifests all the observed degrees of freedom — **quarks and leptons and antiquarks and antileptons**, the observed **gauge vector and scalar fields** — so that nothing in addition is needed to describe observed fields, as well as the cosmological observations.

- ▶ There are **scalar gauge fields** of **two kinds** due to the **two kinds of the Clifford algebra** objects.
- ▶ They offer the explanation for several so far observed phenomena,
 - for the masses of fermions and weak bosons,
 - for the appearance of the matter-antimatter asymmetry in the universe,
 - for the appearance of the dark matter,
 - and for several other phenomena,
- ▶ making several predictions.
- ▶ No assumptions for the group contents of fermions to represent spins, charges and families are needed, what other theories must make.
- ▶ However, **the spontaneous breaks of symmetries are needed.**

A short look "inside" the **spin-charge-family** theory.

The Review article on what is the **spin-charge-family** theory and what are achievements so far is accepted in Progress of Particle and Nuclear Physics.

There are only two kinds of the Clifford algebra objects in any d

- :
- ▶ The **Dirac γ^a operators** (used by Dirac 90 years ago).
 - ▶ The **second one: $\tilde{\gamma}^a$** , which I recognized in the Grassmann space.
 - ▶ **The two kinds form two independent spaces.**

$$\{\gamma^a, \gamma^b\}_+ = 2\eta^{ab} = \{\tilde{\gamma}^a, \tilde{\gamma}^b\}_+,$$

$$\{\gamma^a, \tilde{\gamma}^b\}_+ = 0,$$

The **postulate**

$$(\tilde{\gamma}^a \mathbf{B} = \mathbf{i}(-)^{n_B} \mathbf{B} \gamma^a) |\psi_0\rangle,$$

$$(\mathbf{B} = a_0 + a_a \gamma^a + a_{ab} \gamma^a \gamma^b + \dots + a_{a_1 \dots a_d} \gamma^{a_1} \dots \gamma^{a_d}) |\psi_0\rangle,$$

where $(-)^{n_B} = +1, -1$, if B has a Clifford even or odd character, respectively, and $|\psi_0\rangle$ is a vacuum state on which the operators

γ^a apply,

reduces the Clifford space for the factor of two, while the

operators $\tilde{\gamma}^a \tilde{\gamma}^b$ define the **family quantum numbers**.

$$\mathbf{S}^{ab} := (\mathbf{i}/4)(\gamma^a \gamma^b - \gamma^b \gamma^a),$$

$$\tilde{\mathbf{S}}^{ab} := (\mathbf{i}/4)(\tilde{\gamma}^a \tilde{\gamma}^b - \tilde{\gamma}^b \tilde{\gamma}^a),$$

$$\{\mathbf{S}^{ab}, \tilde{\mathbf{S}}^{cd}\}_- = \mathbf{0}.$$

- ▶ $\tilde{\mathbf{S}}^{ab}$ define the equivalent representations with respect to \mathbf{S}^{ab} .

My recognition:

- ▶ If γ^a are used to describe **the spin and the charges of spinors**,
 $\tilde{\gamma}^a$ - since it must be used or it must be explained why it does not manifest - it **must be used to describe families of spinors**.

Phys. Lett. **B 292**, 25-29 (1992),

J. Math. Phys. **34**, 3731-3745 (1993),

Mod. Phys. Lett. **A 10**, 587-595 (1995)

A simple action for a **spinor** which carries in $d = (13 + 1)$ only **two kinds of spins** (no charges) and for the **gauge gravitational fields**, with which spinors interact:

$$\mathbf{S} = \int d^d x E \mathcal{L}_f + \int d^d x E (\alpha R + \tilde{\alpha} \tilde{R})$$



$$\mathcal{L}_f = \frac{1}{2} (\bar{\psi} \gamma^a p_{0a} \psi) + h.c.$$

$$p_{0a} = f^\alpha{}_a p_{0\alpha} + \frac{1}{2E} \{p_\alpha, E f^\alpha{}_a\} -$$

$$\mathbf{p}_{0\alpha} = \mathbf{p}_\alpha - \frac{1}{2} \mathbf{S}^{ab} \omega_{ab\alpha} - \frac{1}{2} \tilde{\mathbf{S}}^{ab} \tilde{\omega}_{ab\alpha}$$

- ▶ The Einstein action for a free gravitational field is assumed to be linear in the curvature

$$\begin{aligned}
 \mathcal{L}_g &= E (\alpha \mathbf{R} + \tilde{\alpha} \tilde{\mathbf{R}}), \\
 \mathbf{R} &= f^\alpha [a f^{\beta b}] (\omega_{ab\alpha,\beta} - \omega_{ca\alpha} \omega^c_{b\beta}), \\
 \tilde{\mathbf{R}} &= f^\alpha [a f^{\beta b}] (\tilde{\omega}_{ab\alpha,\beta} - \tilde{\omega}_{ca\alpha} \tilde{\omega}^c_{b\beta}),
 \end{aligned}$$

with $E = \det(e^a_\alpha)$
 and $f^\alpha [a f^{\beta b}] = f^{\alpha a} f^{\beta b} - f^{\alpha b} f^{\beta a}$.

- ▶ The internal space of **spinors** (**fermions**) are described by superposition of the odd products of the γ^a 's forming irreducible representations, the family quantum numbers of which are determined by \tilde{S}^{ab} .
- ▶ The only **gauge fields** are the **gravitational** ones – **vielbeins and the two kinds of spin connections**, the gauge fields of S^{ab} and \tilde{S}^{ab} .
- ▶ Both, γ^a and $\tilde{\gamma}^a$, transform as vectors in d ,
and correspondingly also $f^\alpha_a \omega_{bc\alpha}$ and $f^\alpha_a \tilde{\omega}_{bc\alpha}$ transform as tensors with respect to the flat index a .

**

Variation of the action brings for $\omega_{ab\alpha}$

$$\begin{aligned}\omega_{ab\alpha} = & -\frac{1}{2E} \left\{ e_{e\alpha} e_{b\gamma} \partial_\beta (E f^{\gamma[e} f^{\beta]}_a) + e_{e\alpha} e_{a\gamma} \partial_\beta (E f^\gamma_{[b} f^{\beta e]}) \right. \\ & \left. - e_{e\alpha} e^e{}_\gamma \partial_\beta (E f^\gamma_{[a} f^\beta_{b]}) \right\} \\ & - \frac{e_{e\alpha}}{4} \left\{ \bar{\Psi} \left(\gamma_e S_{ab} + \frac{3i}{2} (\delta_b^e \gamma_a - \delta_a^e \gamma_b) \right) \Psi \right\} \\ & - \frac{1}{d-2} \left\{ e_{a\alpha} \left[\frac{1}{E} e^d{}_\gamma \partial_\beta (E f^\gamma_{[d} f^\beta_{b]}) + \frac{1}{2} \bar{\Psi} \gamma^d S_{db} \Psi \right] \right. \\ & \left. - e_{b\alpha} \left[\frac{1}{E} e^d{}_\gamma \partial_\beta (E f^\gamma_{[d} f^\beta_{a]}) + \frac{1}{2} \bar{\Psi} \gamma^d S_{da} \Psi \right] \right\}\end{aligned}$$

IARD, J. Phys.: Conf. Ser. 845 012017 and the refs. therein

**

and for $\tilde{\omega}_{ab\alpha}$,

$$\begin{aligned}\tilde{\omega}_{ab\alpha} = & -\frac{1}{2E} \left\{ e_{e\alpha} e_{b\gamma} \partial_\beta (E f^{\gamma[e} f^{\beta]}_a) + e_{e\alpha} e_{a\gamma} \partial_\beta (E f^{\gamma}{}_{[b} f^{\beta e]}) \right. \\ & \left. - e_{e\alpha} e^e{}_\gamma \partial_\beta (E f^{\gamma}{}_{[a} f^{\beta]}_b) \right\} \\ & - \frac{e_{e\alpha}}{4} \left\{ \bar{\Psi} \left(\gamma_e \tilde{S}_{ab} + \frac{3i}{2} (\delta_b^e \gamma_a - \delta_a^e \gamma_b) \right) \Psi \right\} \\ & - \frac{1}{d-2} \left\{ e_{a\alpha} \left[\frac{1}{E} e^d{}_\gamma \partial_\beta (E f^{\gamma}{}_{[d} f^{\beta]}_b) + \frac{1}{2} \bar{\Psi} \gamma^d \tilde{S}_{db} \Psi \right] \right. \\ & \left. - e_{b\alpha} \left[\frac{1}{E} e^d{}_\gamma \partial_\beta (E f^{\gamma}{}_{[d} f^{\beta]}_a) + \frac{1}{2} \bar{\Psi} \gamma^d \tilde{S}_{da} \Psi \right] \right\}\end{aligned}$$

Eur. Phys. J. C, **77** (2017) 231 and the refs. therein.

If there are no spinors present, the two spin connections are uniquely described by vielbeins $f^\alpha{}_a$.

Fermions

- ▶ The action for **spinors "seen"** from $d = (3 + 1)$ and **analyzed** with respect to the standard model groups as subgroups of $SO(13 + 1)$:

$$\begin{aligned}
 \mathcal{L}_f = & \bar{\psi} \gamma^m (p_m - \sum_{A,i} g^{A\tau} A_i^{Ai}) \psi + \\
 & \{ \sum_{s=[7],[8]} \bar{\psi} \gamma^s p_{0s} \psi \} + \\
 & \{ \sum_{s=[5],[6]} \bar{\psi} \gamma^s p_{0s} \psi + \\
 & \sum_{t=[9],\dots,[14]} \bar{\psi} \gamma^t p_{0t} \psi \} \cdot \\
 & + \text{the rest} , ,
 \end{aligned}$$

Covariant momenta

$$p_{0m} = \left\{ p_m - \sum_A g^A \vec{\tau}^A \vec{A}_m^A \right\}$$

$$\mathbf{m} \quad n \quad (0, 1, 2, 3),$$

$$p_{0s} = f_s^\sigma \left[p_\sigma - \sum_A g^A \vec{\tau}^A \vec{A}_\sigma^A - \sum_A \tilde{g}^A \vec{\tau}^A \vec{A}_\sigma^A \right],$$

$$\mathbf{s} \in (7, 8),$$

$$p_{0s} = f_s^\sigma \left[p_\sigma - \sum_A g^A \vec{\tau}^A \vec{A}_\sigma^A - \sum_A \tilde{g}^A \vec{\tau}^A \vec{A}_\sigma^A \right],$$

$$\mathbf{s} \in (5, 6),$$

$$p_{0t} = f_t^{\sigma'} \left(p_{\sigma'} - \sum_A g^A \vec{\tau}^A \vec{A}_{\sigma'}^A - \sum_A \tilde{g}^A \vec{\tau}^A \vec{A}_{\sigma'}^A \right),$$

$$\mathbf{t} \in (9, 10, 11, \dots, 14),$$

$$\mathbf{A}_s^{\text{Ai}} = \sum_{a,b} \mathbf{c}_{ab}^{\text{Ai}} \omega_{\text{abs}} ,$$

$$\mathbf{A}_t^{\text{Ai}} = \sum_{a,b} \mathbf{c}_{ab}^{\text{Ai}} \omega_{\text{abt}} ,$$

$$\tilde{\mathbf{A}}_s^{\text{Ai}} = \sum_{a,b} \tilde{\mathbf{c}}_{ab}^{\text{Ai}} \tilde{\omega}_{\text{abs}} ,$$

$$\tilde{\mathbf{A}}_t^{\text{Ai}} = \sum_{a,b} \tilde{\mathbf{c}}_{ab}^{\text{Ai}} \tilde{\omega}_{\text{abt}} .$$

$$\tau^{Ai} = \sum_{a,b} c^{Ai}_{ab} \mathbf{S}^{ab},$$

$$\tilde{\tau}^{Ai} = \sum_{a,b} \tilde{c}^{Ai}_{ab} \tilde{\mathbf{S}}^{ab},$$

$$\{\tau^{Ai}, \tau^{Bj}\}_- = i\delta^{AB} f^{Aijk} \tau^{Ak},$$

$$\{\tilde{\tau}^{Ai}, \tilde{\tau}^{Bj}\}_- = i\delta^{AB} f^{Aijk} \tilde{\tau}^{Ak},$$

$$\{\tau^{Ai}, \tilde{\tau}^{Bj}\}_- = 0.$$

- ▶ τ^{Ai} represent the *standard model* charge groups
— $SU(3)_c, SU(2)_w$ — the second $SU(2)_{II}$, the "spinor"
charge $U(1)$, taking care of the hyper charge Y ,
- ▶ $\tilde{\tau}^{Ai}$ denote the family quantum numbers.

$$\mathbf{N}_{(L,R)}^i := \frac{1}{2}(S^{23} \pm iS^{01}, S^{31} \pm iS^{02}, S^{12} \pm iS^{03}),$$

$$\tau_{(1,2)}^i := \frac{1}{2}(S^{58} \mp S^{67}, S^{57} \pm S^{68}, S^{56} \mp S^{78}),$$

$$\tau_3^i := \frac{1}{2}\{S^{9\ 12} - S^{10\ 11}, S^{9\ 11} + S^{10\ 12}, S^{9\ 10} - S^{11\ 12}, \\ S^{9\ 14} - S^{10\ 13}, S^{9\ 13} + S^{10\ 14}, S^{11\ 14} - S^{12\ 13}, \\ S^{11\ 13} + S^{12\ 14}, \frac{1}{\sqrt{3}}(S^{9\ 10} + S^{11\ 12} - 2S^{13\ 14})\},$$

$$\tau^4 := -\frac{1}{3}(S^{9\ 10} + S^{11\ 12} + S^{13\ 14}),$$

$$Y := \tau^4 + \tau^{23},$$

$$Y' := -\tau^4 \tan^2 \vartheta_2 + \tau^{23},$$

$$Q := \tau^{13} + Y,$$

$$Q' := -Y \tan^2 \vartheta_1 + \tau^{13},$$

and equivalently for family groups \tilde{S}^{ab} .

Breaks of symmetries after starting with

- o massless spinors (fermions) ,
- o masses vielbeins and two kinds of the spin connection fields

We prove for a toy model that breaking symmetry in Kaluza-Klein theories can lead to massless fermions.

New J. Phys. 13, 103027, 2011.

J. Phys. A: Math. Theor. 45, 465401, 2012.

[arXiv:1205.1714], [arXiv:1312.541], [arXiv:hep-ph/0412208
p.64-84].

[arXiv:1302.4305], p. 157-166.

$$\text{SO}(1, 13) \times \widetilde{\text{SO}}(1, 13)$$

BREAK I
at $E \geq 10^{16} \text{ GeV}$



$$\text{SO}(1, 7) \times \widetilde{\text{SO}}(1, 7)$$

$$\text{U}(1) \times$$

$$\text{SU}(3)$$



eight massless families



$$\text{SO}(1, 3) \times \text{SO}(4) \times \text{U}(1) \times$$

$$(\widetilde{\text{SU}}(2)_{\widetilde{\text{SO}}(1,3)} \times \widetilde{\text{SU}}(2)_{\widetilde{\text{SO}}(4)}) \times$$

(divided into two groups)

$$(\widetilde{\text{SU}}(2)_{\text{II}\widetilde{\text{SO}}(1,3)} \times \widetilde{\text{SU}}(2)_{\text{II}\widetilde{\text{SO}}(4)}) \times \text{SU}(3)$$

BREAK II



The Standard Model like way of breaking



$$\text{SO}(1, 3) \times \text{U}(1) \times \text{SU}(3)$$

× (two groups of four massive families)

- ▶ Both breaks, the one from $SO(13, 1)$ to $SO(7, 1) \times SO(6)$ and the appearance of the condensate, leave **eight families** ($2^{8/2-1} = 8$, determined by the symmetry of $\widetilde{SO}(1, 7)$) massless. All the families are $\widetilde{SU}(3)$ chargeless. Phys. Rev. D, 80.083534 (2009)
- ▶ The appearance of the **condensate of the two right handed neutrinos**, coupled to **spin 0**, makes the boson gauge fields, with which the condensate interacts, massive. These gauge fields are:
 - All the scalar gauge fields with the space index $s \geq 5$.
 - The vector ($m \leq 3$) gauge fields with the Y' charges — the superposition of $SU(2)_{II}$ and $U(1)_{II}$ charges. J. Phys.: Conf. Ser. 845 (2017) 012017

The **condensate** has spin $S^{12} = 0$, $S^{03} = 0$,
 weak charge $\vec{\tau}^1 = 0$, and
 $\vec{\tilde{\tau}}^1 = 0$, $\tilde{Y} = 0$, $\tilde{Q} = 0$, $\vec{\tilde{N}}_L = 0$.

state	τ^{23}	τ^4	Y	Q	$\tilde{\tau}^{23}$	\tilde{N}_R^3	$\tilde{\tau}^4$
$ \nu_{1R}^{VIII} \rangle_1 \nu_{2R}^{VIII} \rangle_2$	1	-1	0	0	1	1	-1
$ \nu_{1R}^{VIII} \rangle_1 e_{2R}^{VIII} \rangle_2$	0	-1	-1	-1	1	1	-1
$ e_{1R}^{VIII} \rangle_1 e_{2R}^{VIII} \rangle_2$	-1	-1	-2	-2	1	1	-1

- ▶ The **colour, elm, weak and hyper** vector gauge fields do **not interact with the condensate and remain massless.**
J. of Mod. Physics **6** (2015) 2244

- ▶ **At the electroweak break** from $SO(1, 3) \times SU(2)_I \times U(1)_I \times SU(3)$ to $SO(1, 3) \times U(1) \times SU(3)$
 - scalar fields with the space index $s = (7, 8)$ obtain constant values and imaginary masses (nonzero vacuum expectation values,
 - break correspondingly the weak and the hyper charge and change their own masses.
 - They leave massless only the **colour, elm** and **gravity gauge fields**.
- ▶ All the eight massless families gain masses.

Also these is so far just assumed, waiting to be proven that scalar fields, together with boundary conditions, are spontaneously causing also this last breaks.

However, all the needed vector and scalar gauge fields, the fermion fields with all the observed properties, are already in the simple starting action, making the *spin-charge-family* theory (so far) very promising.

- ▶ To the **electroweak break** several scalar fields, the gauge fields of **twice $\widetilde{SU}(2) \times \widetilde{SU}(2)$ and three $\times U(1)$** , contribute, all with the **weak and the hyper charge** of the *standard model Higgs*.
- ▶ They carry besides the **weak** and the **hyper charge** either
 - o the **family members** quantum numbers originating in **(Q,Q',Y')** or
 - o the **family** quantum numbers originating in **twice $\widetilde{SU}(2) \times \widetilde{SU}(2)$** .

J. of Mod. Physics **6** (2015) 2244.



- ▶ The mass matrices of each family member manifest the $\widetilde{SU}(2) \times \widetilde{SU}(2) \times U(1)$ symmetry, which remains unchanged in all loop corrections.

[arXiv:1902.02691, arXiv:1902.10628]

- ▶ We studied on a toy model of $d = (1 + 5)$ conditions which lead after breaking symmetries to massless spinors chirally coupled to the Kaluza-Klein-like gauge field.,

New J. Phys. **13** (2011) 103027, 1-25,

Int. J Mod. Phys. **A 29**, 1450124 (2014), 21 pages.



**

- ▶ There are indeed **two kinds of the Clifford algebra objects**, expressible with Grassmann algebra θ^a 's and their derivatives p^{θ^a} 's ($= i \frac{\partial}{\partial \theta^a}$) in d -dimensional space

$$\gamma^a = (\theta^a - i p^{\theta^a}), \quad \tilde{\gamma}^a = i(\theta^a + i p^{\theta^a}),$$

- ▶ with the properties

$$\{\gamma^a, \gamma^b\}_+ = 2\eta^{ab} = \{\tilde{\gamma}^a, \tilde{\gamma}^b\}_+,$$

$$\{\gamma^a, \tilde{\gamma}^b\}_+ = 0,$$

$$S^{ab} = \frac{i}{4}(\gamma^a \gamma^b - \gamma^b \gamma^a),$$

$$\tilde{S}^{ab} = \frac{i}{4}(\tilde{\gamma}^a \tilde{\gamma}^b - \tilde{\gamma}^b \tilde{\gamma}^a),$$

$$\{S^{ab}, \tilde{S}^{cd}\}_- = 0,$$

manifesting that these two kinds of the Clifford algebra objects, γ^a 's and $\tilde{\gamma}^a$'s form independent algebras (spaces).

► **Postulating**

$$\{\tilde{\gamma}^a \mathbf{B} = (-)^B i \mathbf{B} \gamma^a\} |\psi_{oc}\rangle,$$

with $(-)^B = -1$, if B is (a function of) an odd product of γ^a 's, otherwise $(-)^B = 1$, $|\psi_{oc}\rangle$ is the vacuum state,

- **the operators $\tilde{\gamma}^a$'s offer the definition of the family quantum numbers.**

**Families of quarks and leptons and
antiquarks and antileptons**

Our technique to represent spinors is elegant.

- ▶ J. of Math. Phys. **34**, 3731 (1993),
- ▶ Int. J. of Modern Phys. **A 9**, 1731 (1994),
- ▶ J. of Math. Phys. **36**, 1593 (1995),
- ▶ J. of Math. Phys. **43**, 5782-5803 (2002), hep-th/0111257,
- ▶ J. of Math. Phys. **44** 4817 (2003), hep-th/0303224,
- ▶ J. of High Ener. Phys. **04** (2014) 165, arxiv:1212.2362v2.

The **spinors** states are created out of

nilpotents $(\pm i)^{ab}$ and **projectors** $[\pm i]^{ab}$

$$(\pm i)^{ab} := \frac{1}{2}(\gamma^a \mp \gamma^b), \quad [\pm i]^{ab} := \frac{1}{2}(1 \pm \gamma^a \gamma^b)$$

for $\eta^{aa}\eta^{bb} = -1$,

$$(\pm)^{ab} := \frac{1}{2}(\gamma^a \pm i\gamma^b), \quad [\pm]^{ab} := \frac{1}{2}(1 \pm i\gamma^a \gamma^b),$$

for $\eta^{aa}\eta^{bb} = 1$

with γ^a which are **operators** in d -dimensional space, manifesting Dirac's matrices when applying on "basis vectors" which are odd products of nilpotents and of projectors on vacuum states .

J. of Math. Phys. **34**, 3731 (1993),

Int. J. of Modern Phys. **A 9**, 1731 (1994),

J. of Math. Phys. **43**, 5782-5803 (2002)

Nilpotents $(\pm i)^{ab}$ and **projectors** $[\pm i]^{ab}$ are chosen to be eigensates of the Cartan subalgebra of S^{ab} and \tilde{S}^{ab} .

$$\begin{aligned} S^{ab}(\mathbf{k}) &= \frac{k^{ab}}{2}(\mathbf{k}), & S^{ab}[\mathbf{k}] &= \frac{k^{ab}}{2}[\mathbf{k}], \\ \tilde{S}^{ab}(\mathbf{k}) &= \frac{k^{ab}}{2}(\mathbf{k}), & \tilde{S}^{ab}[\mathbf{k}] &= -\frac{k^{ab}}{2}[\mathbf{k}]. \end{aligned}$$

γ^a transforms $\binom{ab}{k}$ into $\binom{ab}{-k}$, **never** to $\binom{ab}{k}$.

$\tilde{\gamma}^a$ transforms $\binom{ab}{k}$ into $\binom{ab}{k}$, **never** to $\binom{ab}{-k}$.

- ▶ One **Weyl representation** of one **family contains** all the **family members** with the **right handed neutrinos included**. It includes also **antimembers**, reachable by either S^{ab} or by $\mathbb{C}_N \mathcal{P}_N$ on a **family member**.

Jour. of High Energy Phys. **04 (2014)** 165

- ▶ There are $2^{(7+1)/2-1} = 8$ **families**, which decouple into **twice four families**, with the quantum numbers $(\tilde{\tau}^{2i}, \tilde{N}_R^i)$ and $(\tilde{\tau}^{1i}, \tilde{N}_L^i)$, respectively, coupled to corresponding scalar gauge fields.
- ▶ **Besides these two twice triplets coupled to the two groups of family members, all the family members of any of these two groups of family interact with three singlet scalar fields, the gauge fields of τ^4 , Y and Q .**

J. of Math. Phys. **34**, 3731 (1993),

Int. J. of Modern Phys. **A 9**, 1731 (1994),

J. of Math. Phys. **44** 4817 (2003), hep-th/030322.,

S^{ab} generate **all the members of one family**. The **eightplet** (represent. of $SO(7,1)$) of quarks of a particular colour charge

i		$ ^a\psi_i\rangle$	$\Gamma^{(3,1)}$	S^{12}	$\Gamma^{(4)}$	τ^{13}	τ^{23}	Y	τ^4
		Octet, $\Gamma^{(7,1)} = 1$, $\Gamma^{(6)} = -1$, of quarks							
1	u_R^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)(+) & & (+)(+) & & (+) & (-) & (-) \end{matrix}$	1	$\frac{1}{2}$	1	0	$\frac{1}{2}$	$\frac{2}{3}$	$\frac{1}{6}$
2	u_R^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i][-] & & (+)(+) & & (+) & (-) & (-) \end{matrix}$	1	$-\frac{1}{2}$	1	0	$\frac{1}{2}$	$\frac{2}{3}$	$\frac{1}{6}$
3	d_R^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)(+) & & [-][-] & & (+) & (-) & (-) \end{matrix}$	1	$\frac{1}{2}$	1	0	$-\frac{1}{2}$	$-\frac{1}{3}$	$\frac{1}{6}$
4	d_R^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i][-] & & [-][-] & & (+) & (-) & (-) \end{matrix}$	1	$-\frac{1}{2}$	1	0	$-\frac{1}{2}$	$-\frac{1}{3}$	$\frac{1}{6}$
5	d_L^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i](+) & & [-](+) & & (+) & (-) & (-) \end{matrix}$	-1	$\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{6}$	$\frac{1}{6}$
6	d_L^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)[-] & & [-](+) & & (+) & (-) & (-) \end{matrix}$	-1	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{6}$	$\frac{1}{6}$
7	u_L^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i](+) & & (+)[-] & & (+) & (-) & (-) \end{matrix}$	-1	$\frac{1}{2}$	-1	$\frac{1}{2}$	0	$\frac{1}{6}$	$\frac{1}{6}$
8	u_L^{c1}	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)[-] & & (+)[-] & & (+) & (-) & (-) \end{matrix}$	-1	$-\frac{1}{2}$	-1	$\frac{1}{2}$	0	$\frac{1}{6}$	$\frac{1}{6}$

$\gamma^0\gamma^7$ and $\gamma^0\gamma^8$ transform u_R of the 1st row into u_L of the 7th row, and d_R of the 4th row into d_L of the 6th row, doing what the Higgs scalars and γ^0 do in the *standard model*.

The **anti-octet** (repres. of $SO(7,1)$) of **anti-quarks** of the anti-colour charge, reachable by either S^{ab} or $\mathcal{C}_N \mathcal{P}_N^{(d-1)}$:

i		$ ^a \psi_i \rangle$	$\Gamma^{(3,1)}$	S^{12}	$\Gamma^{(4)}$	τ^{13}	τ^{23}	Y	τ^4
		Antioctet, $\Gamma^{(7,1)} = -1$, $\Gamma^{(6)} = 1$, of antiquarks							
33	$\bar{d}_L^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i](+) & & (+)(+) & & [-] & [+] & [+] \end{matrix}$	-1	$\frac{1}{2}$	1	0	$\frac{1}{2}$	$\frac{1}{3}$	$-\frac{1}{6}$
34	$\bar{d}_L^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)[-] & & (+)(+) & & [-] & [+] & [+] \end{matrix}$	-1	$-\frac{1}{2}$	1	0	$\frac{1}{2}$	$\frac{1}{3}$	$-\frac{1}{6}$
35	$\bar{u}_L^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i](+) & & - & & [-] & [+] & [+] \end{matrix}$	-1	$\frac{1}{2}$	1	0	$-\frac{1}{2}$	$-\frac{2}{3}$	$-\frac{1}{6}$
36	$\bar{u}_L^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)[-] & & - & & [-] & [+] & [+] \end{matrix}$	-1	$-\frac{1}{2}$	1	0	$-\frac{1}{2}$	$-\frac{2}{3}$	$-\frac{1}{6}$
37	$\bar{d}_R^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)(+) & & (+)[-] & & [-] & [+] & [+] \end{matrix}$	1	$\frac{1}{2}$	-1	$\frac{1}{2}$	0	$-\frac{1}{6}$	$-\frac{1}{6}$
38	$\bar{d}_R^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i][-] & & (+)[-] & & [-] & [+] & [+] \end{matrix}$	1	$-\frac{1}{2}$	-1	$\frac{1}{2}$	0	$-\frac{1}{6}$	$-\frac{1}{6}$
39	$\bar{u}_R^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ (+i)(+) & & [-](+) & & [-] & [+] & [+] \end{matrix}$	1	$\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$-\frac{1}{6}$	$-\frac{1}{6}$
40	$\bar{u}_R^{\bar{c}1}$	$\begin{matrix} 03 & 12 & 56 & 78 & 9 & 1011 & 1213 & 14 \\ [-i][-] & & [-](+) & & [-] & [+] & [+] \end{matrix}$	1	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$-\frac{1}{6}$	$-\frac{1}{6}$

$\gamma^0 \gamma^7$ and $\gamma^0 \gamma^8$ transform \bar{d}_L of the 1st row into \bar{d}_R of the 5th row, and \bar{u}_L of the 4th row into \bar{u}_R of the 8th row.

New way of second quantization of fermion fields with the internal space described by the odd Clifford algebra manifesting families and family members of quarks and leptons and antiquarks and antileptons

Explaining the Dirac postulates for second quantized fermions.

Any family of the irreducible Lorentz representation of the odd Clifford algebra of $SO(13,1)$ includes **64** members—all the quarks and the leptons, the antiquarks and the antileptons, left and right handed, with spin up and spin down—with the following properties:

- ▶ Quarks distinguish from leptons only in the $SU(3) \times U(1)$ subgroup of the group $SO(13,1)$
- ▶ The $SO(7,1)$ subgroup with **8** elements is the same for quarks and leptons, and the same for antiquarks and antileptons.
- ▶ The Hermitian conjugated representation belong to another odd irreducible representation of the group $SO(13,1)$.

- ▶ Recognizing that the creation operator in the internal space, applying on the vacuum state $|\psi_0\rangle$

$$u_{\uparrow R}^{c1\dagger} \Leftrightarrow b_1^{1\dagger} := (+i)[+ | + || (+)[-][-],$$

- ▶ while its Hermitian conjugated partner is

$$u_{\uparrow R}^{c1} \Leftrightarrow (\mathbf{b}_1^{1\dagger})^\dagger = [-]- || (-)[+ | [+](\mathbf{-i}),$$

- ▶ $u_{\downarrow R}^{c1\dagger} \Leftrightarrow b_1^{2\dagger} := [-i](-) | + || (+)[-][-],$ and

- ▶ $\hat{b}_2^1 := [-]- || (-)[+ | (+)[-i],$

- ▶

- ▶

Creation and Hermitian conjugated partners, operating on a vacuum state

- ▶ $|\psi_0\rangle = [-i][-][-] \cdots [-] |0\rangle$

- ▶ define fermion states—quarks and leptons and antiquarks and antileptons—of one of 8 families.

One finds the following anti-commutation relations for a fermion fields, which represent **"basis vectors" in internal space** determining properties of the observed **quarks and leptons** ($i = (u_{R,L}^{f,\uparrow,\downarrow}, d_{R,L}^{f,\uparrow,\downarrow}, \nu_{R,L}^{f,\uparrow,\downarrow}, e_{R,L}^{f,\uparrow,\downarrow})$) and anti-quarks and anti-leptons , with the family quantum number f .

- ▶ $\{\mathbf{b}_f^m, \mathbf{b}_{f'}^{k\dagger}\}_{*A} |\psi_0\rangle = \delta_{ff'} \delta^{mk} |\psi_0\rangle ,$
- ▶ $\{\mathbf{b}_f^m, \mathbf{b}_{f'}^k\}_+ |\psi_0\rangle = 0 \cdot |\psi_0\rangle ,$
- ▶ $\{\mathbf{b}_f^{m\dagger}, \mathbf{b}_{f'}^{k\dagger}\}_+ |\psi_0\rangle = 0 \cdot |\psi_0\rangle ,$
- ▶ $\mathbf{b}_f^m |\psi_0\rangle = 0 \cdot |\psi_0\rangle ,$
- ▶ $\mathbf{b}_f^{m\dagger} |\psi_0\rangle = |\psi_f^m\rangle$

[arXiv:1802.05554v1], [arXiv:1802.05554v4], [arXiv:1902.10628]

- ▶ There are $2^{\frac{d}{2}-1}$ different "basis vectors" for each of $2^{\frac{d}{2}-1}$ families in d -dimensional space, if there is no break of symmetries.
- ▶ They are defined for any momentum (p^0, \vec{p}) in any dimension d .
- ▶ The "basis vectors" manifest in $d = (3 + 1)$ space spins, charges and families, but not yet include the momentum space.

Tensor products of "basis vectors" of internal space and basis in ordinary space

► Basis in momentum space

$$\begin{aligned} |\vec{p}\rangle &= \hat{b}_{\vec{p}}^\dagger |0_p\rangle, & \langle \vec{p}| &= \langle 0_p| \hat{b}_{\vec{p}}, \\ \langle \vec{p}|\vec{p}'\rangle &= \delta(\vec{p} - \vec{p}') = \langle 0_p| \hat{b}_{\vec{p}} \hat{b}_{\vec{p}'}^\dagger |0_p\rangle, \end{aligned}$$

leading to

$$\hat{b}_{\vec{p}'} \hat{b}_{\vec{p}}^\dagger = \delta(\vec{p}' - \vec{p}),$$

$$\langle 0_p|0_p\rangle = 1, \text{ identity.}$$



$$\begin{aligned} \langle \vec{p}|\vec{x}\rangle &= \langle 0_{\vec{p}}| \hat{b}_{\vec{p}} \hat{b}_{\vec{x}}^\dagger |0_{\vec{x}}\rangle = (\langle 0_{\vec{x}}| \hat{b}_{\vec{x}} \hat{b}_{\vec{p}}^\dagger |0_{\vec{p}}\rangle)^\dagger \\ \{\hat{b}_{\vec{p}}^\dagger, \hat{b}_{\vec{p}'}^\dagger\}_- &= 0, & \{\hat{b}_{\vec{p}}, \hat{b}_{\vec{p}'}\}_- &= 0, & \{\hat{b}_{\vec{p}}, \hat{b}_{\vec{p}'}^\dagger\}_- &= 0, \\ \{\hat{b}_{\vec{x}}^\dagger, \hat{b}_{\vec{x}'}^\dagger\}_- &= 0, & \{\hat{b}_{\vec{x}}, \hat{b}_{\vec{x}'}\}_- &= 0, & \{\hat{b}_{\vec{x}}, \hat{b}_{\vec{x}'}^\dagger\}_- &= 0, \end{aligned}$$

while

$$\{\hat{b}_{\vec{p}}, \hat{b}_{\vec{x}}^\dagger\}_- = e^{i\vec{p}\cdot\vec{x}} \frac{1}{\sqrt{(2\pi)^{d-1}}}, \quad \{\hat{b}_{\vec{x}}, \hat{b}_{\vec{p}}^\dagger\}_- = e^{-i\vec{p}\cdot\vec{x}} \frac{1}{\sqrt{(2\pi)^{d-1}}},$$

► **Creation and annihilation operators in internal and ordinary space**

$$\{\hat{\mathbf{b}}_f^{s\dagger}(\vec{p})|_{p^0=|\vec{p}|} = \sum_m c^{ms}_f(\vec{p}, p^0=|\vec{p}|) \hat{b}_{\vec{p}}^{\dagger *T} \hat{b}_f^{m\dagger}\} |\psi_o\rangle = |0_{\vec{p}}\rangle,$$

with the coefficients $c^{ms}_f(\vec{p}, p^0=|\vec{p}|)$ chosen so, that $\hat{\mathbf{b}}_f^{s\dagger}(\vec{p})|_{p^0=|\vec{p}|} \cdot e^{-ip_a x^a}$ solve, applied on the vacuum state, the Weyl equations for **free massless fermions**.



$$\mathcal{A} = \int d^d x \frac{1}{2} (\psi^\dagger \gamma^0 \gamma^a \mathbf{p}_a \psi) + \text{h.c.},$$

$$\gamma^a \mathbf{p}_a |\psi\rangle = \mathbf{0},$$

$$|\psi_f^s(\vec{p})\rangle = \hat{\mathbf{b}}_f^{s\dagger}(\vec{p})|_{p^0=|\vec{p}|} |\psi_o\rangle = |0_{\vec{p}}\rangle,$$

- ▶ The commutation relations for $\hat{\mathbf{b}}_f^s(\vec{p})$ and $\hat{\mathbf{b}}_f^{s\dagger}(\vec{p})$

$$\{\hat{\mathbf{b}}_{f'}^{s'}(\vec{p}'), \hat{\mathbf{b}}_f^{s\dagger}(\vec{p})\}_+ |\psi_{oc}\rangle |0_{\vec{p}}\rangle = \delta^{ss'} \delta_{ff'} \delta(\vec{p}' - \vec{p}) |\psi_{oc}\rangle |0_{\vec{p}}\rangle$$

$$\{\hat{\mathbf{b}}_{f'}^{s'}(\vec{p}'), \hat{\mathbf{b}}_f^s(\vec{p})\}_+ |\psi_{oc}\rangle |0_{\vec{p}}\rangle = 0 |\psi_{oc}\rangle |0_{\vec{p}}\rangle,$$

$$\{\hat{\mathbf{b}}_{f'}^{s'\dagger}(\vec{p}'), \hat{\mathbf{b}}_f^{s\dagger}(\vec{p})\}_+ |\psi_{oc}\rangle |0_{\vec{p}}\rangle = 0 |\psi_{oc}\rangle |0_{\vec{p}}\rangle,$$

$$\hat{\mathbf{b}}_f^{s\dagger}(\vec{p}) |\psi_{oc}\rangle |0_{\vec{p}}\rangle = |\psi_f^s(\vec{p})\rangle$$

$$\hat{\mathbf{b}}_f^s(\vec{p}) |\psi_{oc}\rangle |0_{\vec{p}}\rangle = 0 |\psi_{oc}\rangle |0_{\vec{p}}\rangle$$

$$p^0 = |\vec{p}|.$$

- ▶ **Creation and annihilation operators, $\hat{\mathbf{b}}_f^{s\dagger}(\vec{p})$ and $\hat{\mathbf{b}}_f^s(\vec{p})$, fulfill anticommutation relations due to the Clifford oddness of the "basis vectors" describing the internal space of fermions.**
- ▶ **No postulates of Dirac kind are needed.**
- ▶ **Describing the internal space with the odd Clifford algebra "basis vectors" explains correspondingly the Dirac postulates.**

- ▶ One only needs to prove that the commutational relations are valid also on the whole Hilbert space of all possible products of all possible creation operators on the vacuum state.
- ▶ We have for each momentum $\tilde{\mathbf{p}}$ correspondingly tensor products of one single particle state, of two single particle states, up to $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ **single particle states**, if there is no break of symmetry.

- ▶ A tensor product of $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ single particle states of all possible (∞ many) momenta \vec{p} , $|p^0| = |\vec{p}|$, form the Hilbert space of all possible Slater determinants with no state occupied, with one state — any one — occupied and all the rest unoccupied, with two states occupied, up to all the $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ states occupied.
- ▶ The Hilbert space $\mathcal{H}_{\vec{p}}$ of a particular momentum \vec{p} consists correspondingly of 2^{2^d-2} “Slater determinants”, since the “Slater determinant” can have no occupied state, one occupied state, two occupied states, up to 2^{d-2} occupied states, respectively.

- ▶ **The total Hilbert space \mathcal{H} is the continuously infinite product $(\otimes_{\mathbf{N}})$ of $\mathcal{H}_{\tilde{\mathbf{p}}}$ with $2^{2^{d-2}}$ “Slater determinants”:**

$$\mathcal{H} = \prod_{\tilde{\mathbf{p}}=-\infty}^{\infty} \otimes_{\mathbf{N}} \mathcal{H}_{\tilde{\mathbf{p}}},$$

where the product $(\otimes_{\mathbf{N}})$ is to point out that Clifford odd objects $\hat{\mathbf{b}}_{\mathbf{f}}^{s\dagger}$ and $\hat{\mathbf{b}}_{\mathbf{f}'}^{s'}$ **“can not forget” that they are Clifford odd objects, no matter to which $\tilde{\mathbf{p}}$ their superposition $\hat{\mathbf{b}}_{\mathbf{f}}^{s\dagger}(\tilde{\mathbf{p}})$, $\mathbf{p}^0 = |\tilde{\mathbf{p}}|$, (solutions of equation of motion) belong.**

- ▶ **The Clifford odd creation and annihilation operators fulfill the anticommutation relations on the whole Hilbert space**

$$\{\hat{\mathbf{b}}_f^{s\dagger}(\vec{\mathbf{p}}), \hat{\mathbf{b}}_{f'}^{s'\dagger}(\vec{\mathbf{p}}')\}_{+\mathcal{H}} = \mathbf{0},$$

$$\{\hat{\mathbf{b}}_f^{s\dagger}(\vec{\mathbf{p}}), \hat{\mathbf{b}}_{f'}^{s'}(\vec{\mathbf{p}}')\}_{+\mathcal{H}} = \mathbf{0},$$

$$\{\hat{\mathbf{b}}_f^{s\dagger}(\vec{\mathbf{p}}), \hat{\mathbf{b}}_{f'}^{s'}(\vec{\mathbf{p}}')\}_{+\mathcal{H}} = \delta^{ss'} \delta_{ff'} \delta(\vec{\mathbf{p}} - \vec{\mathbf{p}}') \mathcal{H},$$

for any of continuously infinite $\vec{\mathbf{p}}$ and any $\vec{\mathbf{p}}'$.

- We can represent $\mathcal{H}_{\tilde{p}}$ as follows

$$\begin{aligned}
|0_{1p}, 0_{2p}, 0_{3p}, \dots, 0_{2^{d-2}p} > |1 & , \\
|1_{1p}, 0_{2p}, 0_{3p}, \dots, 0_{2^{d-2}p} > |2 & , \\
|0_{1p}, 1_{2p}, 0_{3p}, \dots, 0_{2^{d-2}p} > |3 & , \\
|0_{1p}, 0_{2p}, 1_{3p}, \dots, 0_{2^{d-2}p} > |4 & , \\
& \vdots \\
|1_{1p}, 1_{2p}, 0_{3p}, \dots, 0_{2^{d-2}p} > |2^{d-2}+2 & , \\
& \vdots \\
|1_{1p}, 1_{2p}, 1_{3p}, \dots, 1_{2^{d-2}p} > |2^{2^{d-2}} & .
\end{aligned}$$

repeating these Slater determinants for all ∞ number of \tilde{p} .

Let us compare the above **second quantization procedure**, describing internal space of fermions with the odd Clifford algebra objects, with the **second quantization procedure proposed by Dirac**, where the creation operators and correspondingly their Hermitian conjugate operators are **assumed**.

$$\psi_i(\mathbf{t}, \tilde{\mathbf{x}}) = \sum_{\mathbf{p}, i} \hat{a}^\dagger(\mathbf{p}, i) \mathbf{v}(\tilde{\mathbf{p}}, i) e^{-\mathbf{p}_a \mathbf{x}^a}.$$

$\mathbf{v}(\tilde{\mathbf{p}}, i)$ determine solutions of equations of motion for a particular $e^{-\mathbf{p}_a \mathbf{x}^a}$.

$\hat{a}^\dagger(p, i)$ is just assumed, together with the (assumed) Hermitian conjugate operator, to fill the anticommutation relation.

In the spin-charge-family theory the creation operators appear from the odd Clifford objects, representing fermion states, applying on the vacuum state, in internal space.

- ▶ The **anticommutation relations for creation operators and their Hermitian conjugated partners in the Dirac case** in $d = (3 + 1)$ for **spin** (\uparrow, \downarrow) and **right and left handedness** (± 1 , respectively)

$$\begin{aligned} \{\hat{a}_i^\dagger(\vec{p}), \hat{a}_j^\dagger(\vec{p}')\}_+ &= 0 = \{\hat{a}_i(\vec{p}), \hat{a}_j(\vec{p}')\}_+, \\ \{\hat{a}_i(\vec{p}), \hat{a}_j^\dagger(\vec{p}')\}_+ &= \delta_{ij} \delta(\vec{p} - \vec{p}'), \end{aligned}$$

in the case of massless fermions.

- ▶ To be able to compare the spin-charge-family theory creation operators for this particular case of $d = (3 + 1)$, we make a choice of the creation operators representing spin \uparrow and right handedness,

$$\hat{b}_1^\dagger := \begin{matrix} 03 & 12 \\ [+i] & (+) \end{matrix},$$

and spin \downarrow and right handedness,

$$\hat{b}_2^\dagger := \begin{matrix} 03 & 12 \\ (-i) & [-] \end{matrix},$$

and shall not pay attention on charges (which in spin-charge-family theory originate in $d \geq 5$) and families.

- ▶ In the spin-charge-family case the creation operators $\hat{b}_i^\dagger, i = (1, 2)$ originate in "basis vectors" determining the internal space of fermions.
- ▶ Solutions of the Weyl equation – the Dirac equation for massless fermions – are superposition of both "basis vectors" for particular $\vec{p}, p^0 = |\vec{p}|$
 $\hat{\mathbf{b}}^{s\dagger}(\vec{p}) = \sum_i c^{is}(\vec{p}) \hat{b}_p^\dagger *_{\mathcal{T}} \hat{b}^{i\dagger} \quad i = (1, 2).$
- ▶ Let us write down both kinds of creation operators, the Dirac one and ours, both for the right handed case, leaving out therefore the index describing handedness h in the Dirac case and f , describing family, in our case

$$\hat{\mathbf{a}}^{s\dagger}(\vec{p}) \stackrel{\text{def}}{=} \sum_i \hat{\mathbf{a}}_i^{s\dagger}(\vec{p}) u_i^s(\vec{p}), \quad \hat{\mathbf{b}}^{s\dagger}(\vec{p}) = \sum_i c^{is}(\vec{p}) \hat{b}^{i\dagger} \hat{b}_p,$$

what is my redefinition of Dirac's operators.

- ▶ The creation operator of Dirac

$$\hat{\mathbf{a}}^{s\dagger}(\vec{p}) = \sum_i u_i^s(\vec{p}) \hat{a}_i^{s\dagger}(\vec{p}), \quad v^{si}(\vec{p}, \vec{x}) = u^{si}(\vec{p}) e^{i\vec{p} \cdot \vec{x}},$$

has to be related to

$$\hat{\mathbf{b}}^{s\dagger} = \sum_i c^{is}(\vec{p}) \hat{b}^i \hat{b}_{\vec{p}}.$$

$$\hat{\mathbf{a}}^{s\dagger}(\vec{p}) = \sum_i \hat{a}_i^{s\dagger}(\vec{p}) u_i^s(\vec{p}) \text{ to be related to } \hat{\mathbf{b}}^{s\dagger}(\vec{p}) = \sum_i c^{is}(\vec{p}) \hat{b}^i \hat{b}_{\vec{p}}.$$

Both creation operators, $\hat{\mathbf{a}}^{s\dagger}(\vec{p})$ and $\hat{\mathbf{b}}^{s\dagger}(\vec{p})$, fulfill the same anticommutation relations,

$\hat{\mathbf{a}}^{s\dagger}(\vec{p})$ fulfill also the anticommutation relations of Dirac.

- ▶ Dirac equipped the creation operators (and correspondingly also the annihilation operators) with the quantum numbers (s, i) and with \vec{p} . He postulated for such creation and annihilation operators anticommutation relations.
- ▶ Our creation and annihilation operators, $\hat{\mathbf{b}}^{s\dagger}(\vec{p})$ and $\hat{\mathbf{b}}^s(\vec{p})$, have anticommuting properties due to the anticommutativity of $\hat{\mathbf{b}}^{i\dagger}$ and $\hat{\mathbf{b}}^i$, which are Clifford odd objects.

The odd Clifford algebra offers the explanation for the Dirac's postulates for the second quantized fermions.

▶ $\psi^s(\vec{x}) = \int_{-\infty}^{+\infty} d^3p \hat{b}^{s\dagger}(\vec{p}) e^{-i(p^0 x^0 - \vec{p}_k \cdot \vec{x})}$

to be related to

$$\psi^s(\vec{x}) = \int_{-\infty}^{+\infty} d^3p \hat{a}^{s\dagger}(\vec{p}) e^{-i(p^0 x^0 - \vec{p}_k \cdot \vec{x})}.$$

The theory explains:

- ▶ The appearance of the **finite** number of the **internal space basis vectors** of fermions.
- ▶ The **anticommutation relations** among the creation and annihilation operators, creating the **anticommuting** single fermion states.
- ▶ The continuously **infinite** number of creation operators due to infinite dimensional ordinary space.
- ▶ The tensor products of **Clifford odd creation operators** explain the **Hilbert space of the second quantized fermions**.

- ▶ It is worthwhile to notice that "nature could make a choice" of Grassmann rather than Clifford space:
 - Also in Grassmann space, namely, one finds the anticommutation relations needed for a fermion field.
 - But in this case spinors would have spins and charges in adjoint representations with respect to particular subgroups.
 - And no families would appear.

Vector gauge fields origin in gravity,
in vielbeins and two kinds of the spin connection fields,
the gauge fields of S^{ab} and \tilde{S}^{ab}

**

- ▶ **All the vector gauge fields**, A_m^{Ai} , $(m, n) = (0, 1, 2, 3)$ of the observed charges $\tau^{Ai} = \sum_{s,t} c^{Aist} S^{st}$, manifesting at the observable energies, **have all the properties as assumed by the standard model**.
- ▶ They carry with respect to the space index $m \in (0, 1, 2, 3)$ the vector degrees of freedom, while they have additional **internal degrees of freedom** (τ^{Ai}) in the adjoint representations.
- ▶ They origin as spin conection gauge fields of S^{ab} :
 $A_m^{Ai} = \sum_{s,t} c^{Aist} \omega_{stm}$.
- ▶ S^{ab} applies on indexes (s, t, m) as follows

$$S^{ab} \omega_{stm\dots g} = i(\delta_s^a \omega_{tm\dots g}^b - \delta_s^b \omega_{tm\dots g}^a).$$

**

The action for vectors with respect to the space index
 $m = (0, 1, 2, 3)$ origin in **gravity**

$$\int \mathbf{E} d^4 \mathbf{x} d^{(d-4)} \mathbf{x} \alpha \mathbf{R}^{(d)} = \int d^4 \mathbf{x} \left\{ -\frac{1}{4} \mathbf{F}^{\mathbf{A}i}{}_{\mathbf{m}n} \mathbf{F}^{\mathbf{A}i\mathbf{m}n} \right\},$$
$$\mathbf{A}^{\mathbf{A}i}{}_{\mathbf{m}} = \sum_{s,t} \mathbf{c}^{\mathbf{A}ist} \omega_{st\mathbf{m}}.$$

Eur. Phys. J. C. **77** (2017) 231,

*

Also scalar fields
(there are doublets and triplets)
origin in gravity fields — **they are spin connections and vielbeins** —
with the space index $s \geq 5$.

Eur. Phys. J. C. **77** (2017) 231 ,
Phys. Rev. **D 91** (2015) 6, 065004,
J. of Mod. Physics **6** (2015) 2244.

- ▶ There are several **scalar gauge fields** with the space index $(s,t,s') = (7,8)$, all origin in the spin connection fields, either $\tilde{\omega}_{abs}$ or $\omega_{s'ts}$:
 - Twice **two triplets**, the scalar gauge fields with the **family** quantum numbers $(\tilde{\tau}^{Ai} = \sum_{a,b} \tilde{c}^{Ai}_{ab} \tilde{S}^{ab})$ and
 - **three singlets** with the **family members** quantum numbers (Q,Q',Y') , the gauge fields of S^{st} .
- ▶ They are all doublets with respect to the space index **(5,6,7,8)**.
- ▶ They have all the rest quantum numbers **determined by the adjoint representations**.
- ▶ They explain at the so far observable energies the **Higgs's scalar** and the **Yukawa couplings**.

The two doublets, determining the properties of the Higgs's scalar and the Yukawa couplings, are:

	state	τ^{13}	$\tau^{23} = Y$	spin	τ^4	Q
A_{78}^{Ai} (-)	$A_7^{Ai} + iA_8^{Ai}$	$+\frac{1}{2}$	$-\frac{1}{2}$	0	0	0
A_{56}^{Ai} (-)	$A_5^{Ai} + iA_6^{Ai}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	0	-1
A_{78}^{Ai} (+)	$A_7^{Ai} - iA_8^{Ai}$	$-\frac{1}{2}$	$+\frac{1}{2}$	0	0	0
A_{56}^{Ai} (+)	$A_5^{Ai} - iA_6^{Ai}$	$+\frac{1}{2}$	$+\frac{1}{2}$	0	0	+1

There are A_{78}^{Ai} and A_{78}^{Ai} which gain **nonzero vacuum expectation values** at the **electroweak break**.

Index Ai determines the **family** ($\tilde{\tau}^{Ai}$) quantum numbers and the **family members** (Q,Q',Y') quantum numbers, both are in adjoint representations.

- ▶ There are besides **doublets**, with the space index $s = (5, 6, 7, 8)$, as well **triplets** and **anti-triplets**, with respect to the space index $s = (9, \dots, 14)$.
- ▶ **There are no additional scalars** in the theory for **$d=(13+1)$** .
- ▶ **All are massless.**
- ▶ All the scalars have the family and the family members quantum numbers in the **adjoint** representations.
- ▶ The properties of scalars are to be analyzed with respect to the generators of the corresponding subgroups, expressible with S^{ab} , as it is the case of the vector gauge fields.
- ▶ It is the (**so far assumed**) **condensate**, which makes those gauge fields, with which it interacts, massive.
 - o **The condensate breaks the CP symmetry.**

- ▶ The **scalar condensate** of two **right handed neutrinos** couple to
 - all the **scalar and vector** gauge fields, making them massive,
 - It does not interact with the **weak charge $SU(2)_I$** , the **hyper charge $U(1)$** , and the **colour $SU(3)$ charge gauge fields**, as well as the **gravity**, leaving them **massless**.

J. of Mod.Phys.**4** (2013) 823-847,

J. of Mod.Phys. **6** (2015) 2244-2247,

Phys Rev.**D 91**(2015)6,065004.

Scalars with $s=(7,8)$, which gain **nonzero vacuum expectation values**, break the **weak and the hyper** symmetry, while conserving the **electromagnetic and colour** charge:

$$\begin{aligned}
 \mathbf{A}_s^{\mathbf{A}i} &\supset (\mathbf{A}_s^{\mathbf{Q}}, \mathbf{A}_s^{\mathbf{Q}'}, \mathbf{A}_s^{\mathbf{Y}'}, \tilde{\tilde{\mathbf{A}}}_s^{\tilde{\tilde{1}}}, \tilde{\tilde{\mathbf{A}}}_s^{\tilde{\tilde{N}}_L}, \tilde{\tilde{\mathbf{A}}}_s^{\tilde{\tilde{2}}}, \tilde{\tilde{\mathbf{A}}}_s^{\tilde{\tilde{N}}_R}), \\
 \tau^{\mathbf{A}i} &\supset (\mathbf{Q}, \mathbf{Q}', \mathbf{Y}', \tilde{\tilde{\tau}}^1, \tilde{\tilde{\mathbf{N}}}_L, \tilde{\tilde{\tau}}^2, \tilde{\tilde{\mathbf{N}}}_R), \\
 \mathbf{s} &= (7, 8).
 \end{aligned}$$

Ai denotes:

o family quantum numbers

$(\tilde{\tilde{\tau}}^1, \tilde{\tilde{\mathbf{N}}}_L)$ quantum numbers of the first group of four families

and

$(\tilde{\tilde{\tau}}^2, \tilde{\tilde{\mathbf{N}}}_R)$ quantum numbers of the second group of four families.

o And family members quantum numbers $(\mathbf{Q}, \mathbf{Q}', \mathbf{Y}')$

A_s^{Ai} are expressible with either $\omega_{sts'}$ or $\tilde{\omega}_{abs'}$.

$$\vec{A}_s^1 = (\tilde{\omega}_{58s} - \tilde{\omega}_{67s}, \tilde{\omega}_{57s} + \tilde{\omega}_{68s}, \tilde{\omega}_{56s} - \tilde{\omega}_{78s}),$$

$$\vec{A}_s^2 = (\tilde{\omega}_{58s} + \tilde{\omega}_{67s}, \tilde{\omega}_{57s} - \tilde{\omega}_{68s}, \tilde{\omega}_{56s} + \tilde{\omega}_{78s}),$$

$$\vec{A}_{Ls}^N = (\tilde{\omega}_{23s} + i\tilde{\omega}_{01s}, \tilde{\omega}_{31s} + i\tilde{\omega}_{02s}, \tilde{\omega}_{12s} + \tilde{\omega}_{03s}),$$

$$\vec{A}_{Rs}^N = (\tilde{\omega}_{23s} - i\tilde{\omega}_{01s}, \tilde{\omega}_{31s} - i\tilde{\omega}_{02s}, \tilde{\omega}_{12s} - i\tilde{\omega}_{03s}),$$

$$A_s^Q = \omega_{56s} - (\omega_{910s} + \omega_{1112s} + \omega_{1314s}),$$

$$A_s^Y = (\omega_{56s} + \omega_{78s}) - (\omega_{910s} + \omega_{1112s} + \omega_{1314s})$$

$$A_s^4 = -(\omega_{910s} + \omega_{1112s} + \omega_{1314s}).$$

The **mass term**, appearing in the **starting action**,
 is (p_s , when treating the lowest energy solutions, is left out)

$$\mathcal{L}_M = \sum_{s=(7,8), Ai} \bar{\psi} \gamma^s (-\tau^{Ai} A_s^{Ai}) \psi =$$

$$-\bar{\psi} \left\{ \overset{78}{(+)} \tau^{Ai} (A_7^{Ai} - i A_8^{Ai}) + \overset{78}{(-)} \tau^{Ai} (A_7^{Ai} + i A_8^{Ai}) \right\} \psi ,$$

$$\overset{78}{(\pm)} = \frac{1}{2} (\gamma^7 \pm i \gamma^8), \quad A_{\overset{78}{(\pm)}}^{Ai} := (A_7^{Ai} \mp i A_8^{Ai}).$$

Operators Y , Q and τ^{13} , applied on $(A_7^{A_i} \mp i A_8^{A_i})$

$$\tau^{13} (A_7^{A_i} \mp i A_8^{A_i}) = \pm \frac{1}{2} (A_7^{A_i} \mp i A_8^{A_i}),$$

$$Y (A_7^{A_i} \mp i A_8^{A_i}) = \mp \frac{1}{2} (A_7^{A_i} \mp i A_8^{A_i}),$$

$$Q (A_7^{A_i} \mp i A_8^{A_i}) = 0,$$

manifest that **all** $(A_7^{A_i} \mp i A_8^{A_i})$ have quantum numbers of the **Higgs's scalar of the standard model**, "dressing", after **gaining nonzero expectation values**, the right handed members of a family with appropriate charges, so that they gain charges of the left handed partners:

$(A_7^{A_i} + i A_8^{A_i})$ "dresses" u_R, ν_R and $(A_7^{A_i} - i A_8^{A_i})$ "dresses" d_R, e_R , with quantum numbers of their left handed partners, just as required by the "standard model".

Ai measures:

either

o the **Q,Q',Y' charges** of the **family members**

or

o **family charges** ($\vec{\tau}^{\vec{I}}, \vec{N}_L$), transforming a family member of one family into the same family member of another family,

manifesting in each group of four families the

$$\widetilde{SU}(2) \times \widetilde{SU}(2) \times U(1)$$

symmetry.

**** Eight families** of u_R (spin 1/2, colour $(\frac{1}{2}, \frac{1}{2\sqrt{3}})$) and of colourless ν_R (spin 1/2). All have "tilde spinor charge" $\tilde{\tau}^4 = -\frac{1}{2}$, the weak charge $\tau^{13} = 0$, $\tau^{23} = \frac{1}{2}$. Quarks have "spinor" q.no. $\tau^4 = \frac{1}{6}$ and leptons $\tau^4 = -\frac{1}{2}$. The first four families have $\tilde{\tau}^{23} = 0$, $\tilde{N}_R^3 = 0$, the second four families have $\tilde{\tau}^{13} = 0$, $\tilde{N}_L^3 = 0$.

$\tilde{N}_R^3 = 0, \tilde{\tau}^{23} = 0$		$\tilde{N}_R^3 = 0, \tilde{\tau}^{23} = 0$		$\tilde{\tau}^{13}$	\tilde{N}_L^3
u_{R1}^{c1}	03 12 56 78 9 10 11 12 13 14 (+i) [+] [+] (+) (+) [-] [-]	ν_{R1}	03 12 56 78 9 10 11 12 13 14 (+i) [+] [+] (+) (+) (+) (+)	$-\frac{1}{2}$	$-\frac{1}{2}$
u_{R2}^{c1}	03 12 56 78 9 10 11 12 13 14 [+ i] (+) [+] (+) (+) [-] [-]	ν_{R2}	03 12 56 78 9 10 11 12 13 14 [+ i] (+) [+] (+) (+) (+) (+)	$-\frac{1}{2}$	$\frac{1}{2}$
u_{R3}^{c1}	03 12 56 78 9 10 11 12 13 14 (+i) [+] (+) [+] (+) [-] [-]	ν_{R3}	03 12 56 78 9 10 11 12 13 14 (+i) [+] (+) [+] (+) (+) (+)	$\frac{1}{2}$	$-\frac{1}{2}$
u_{R4}^{c1}	03 12 56 78 9 10 11 12 13 14 [+ i] (+) (+) [+] (+) [-] [-]	ν_{R4}	03 12 56 78 9 10 11 12 13 14 [+ i] (+) (+) [+] (+) (+) (+)	$\frac{1}{2}$	$\frac{1}{2}$
$\tilde{N}_L^3 = 0, \tilde{\tau}^{13} = 0$		$\tilde{N}_L^3 = 0, \tilde{\tau}^{13} = 0$		$\tilde{\tau}^{23}$	\tilde{N}_R^3
u_{R5}^{c1}	03 12 56 78 9 10 11 12 13 14 (+i) (+) (+) (+) (+) [-] [-]	ν_{R5}	03 12 56 78 9 10 11 12 13 14 (+i) (+) (+) (+) (+) (+) (+)	$-\frac{1}{2}$	$-\frac{1}{2}$
u_{R6}^{c1}	03 12 56 78 9 10 11 12 13 14 (+i) (+) [+] [+] (+) [-] [-]	ν_{R6}	03 12 56 78 9 10 11 12 13 14 (+i) (+) [+] [+] (+) (+) (+)	$-\frac{1}{2}$	$\frac{1}{2}$
u_{R7}^{c1}	03 12 56 78 9 10 11 12 13 14 [+ i] [+] (+) (+) (+) [-] [-]	ν_{R7}	03 12 56 78 9 10 11 12 13 14 [+ i] [+] (+) (+) (+) (+) (+)	$\frac{1}{2}$	$-\frac{1}{2}$
u_{R8}^{c1}	03 12 56 78 9 10 11 12 13 14 [+ i] [+] [+] [+] (+) [-] [-]	ν_{R8}	03 12 56 78 9 10 11 12 13 14 [+ i] [+] [+] [+] (+) (+) (+)	$\frac{1}{2}$	$\frac{1}{2}$

Before the **electroweak break** all the **families** are **mass protected** and correspondingly **massless**.

- ▶ Scalars with the weak and the hyper charge $(\mp\frac{1}{2}, \pm\frac{1}{2})$ determine masses of **all the family members** α of the **lower four families**, ν_R of the lower four families have nonzero $Y' := -\tau^4 + \tau^{23}$ and interact with the scalar field $(A_{(\pm)}^{Y'}, \vec{A}_{(\pm)}^{\tilde{1}}, \vec{A}_{(\pm)}^{\tilde{N}_L})$.
- ▶ The group of the lower four families manifest the $\widetilde{SU}(2)_{\widetilde{SO}(1,3)} \times \widetilde{SU}(2)_{\widetilde{SO}(4)} \times U(1)$ **symmetry** (also after all loop corrections).

$$\mathcal{M}^\alpha = \begin{pmatrix} -a_1 - a & e & d & b \\ e^* & -a_2 - a & b & d \\ d^* & b^* & a_2 - a & e \\ b^* & d^* & e^* & a_1 - a \end{pmatrix}^\alpha.$$

[arXiv:1412.5866], [arXiv:1902.02691], [arXiv:1902.10628]

We **made calculations**, treating **quarks** and **leptons** in equivalent way, as required by the "spin-charge-family" theory. Although

- ▶ any **$(n-1) \times (n-1)$** submatrix of an unitary **$n \times n$** matrix determines the **$n \times n$** matrix for **$n \geq 4$** uniquely,
- ▶ the **measured mixing matrix elements** of the **3×3** submatrix **are not yet accurate enough even for quarks to predict the masses m_4 of the fourth family members.**
 - We can say, taking into account the data for the mixing matrices and masses, that **m_4 quark masses might be any in the interval $(300 < m_4 < 1000)$ GeV or even **above**.** Other experiments require that m_4 are above 1000 GeV.
- ▶ **Assuming** masses **m_4** we can predict mixing matrices.

Results are presented for two choices of $m_{u_4} = m_{d_4}$, [arxiv:1412.5866]:

- ▶ 1. $m_{u_4} = 700$ GeV, $m_{d_4} = 700$ GeV.....new₁
- ▶ 2. $m_{u_4} = 1200$ GeV, $m_{d_4} = 1200$ GeV.....new₂

exp_n	0.97425 ± 0.00022	0.2253 ± 0.0008	0.00413 ± 0.00049	
new ₁	0.97423(4)	0.22539(7)	0.00299	0.00776(1)
new ₂	0.97423[5]	0.22538[42]	0.00299	0.00793[466]
exp_n	0.225 ± 0.008	0.986 ± 0.016	0.0411 ± 0.0013	
new ₁	0.22534(3)	0.97335	0.04245(6)	0.00349(60)
new ₂	0.22531[5]	0.97336[5]	0.04248	0.00002[216]
exp_n	0.0084 ± 0.0006	0.0400 ± 0.0027	1.021 ± 0.032	
new ₁	0.00667(6)	0.04203(4)	0.99909	0.00038
new ₂	0.00667	0.04206[5]	0.99909	0.00024[21]
new ₁	0.00677(60)	0.00517(26)	0.00020	0.99996
new ₂	0.00773	0.00178	0.00022	0.99997[9]

One can see what

B. Belfatto, R. Beradze, Z. Berezhiani, required in [arXiv:1906.02714v1], that

$V_{u_1 d_4} > V_{u_1 d_3}$, $V_{u_2 d_4} < V_{u_1 d_4}$, and $V_{u_3 d_4} < V_{u_1 d_4}$, ,
what is just happening in my theory.

The newest experimental data, PDG, (P A Zyla et al, Prog. Theor. and Exp. Phys., Vol. 2020, Issue 8, Aug. 2020, 083C01) have not yet been used to fit mass matrix of Eq. (2).

- ▶ **o** The **matrix elements** V_{CKM} **depend strongly on the accuracy** of the experimental **3 x 3 submatrix**.
- o** Calculated **3 x 3 submatrix** of 4×4 V_{CKM} depends on the m_{4th} **family masses**, but not much.
- o** $V_{u_i d_4}$, $V_{d_j u_4}$ do not depend strongly on the m_{4th} family masses and are obviously **very small**.
- ▶ The higher are the fourth family members masses, the closer are the mass matrices to the **democratic matrices** for either quarks or leptons, as expected.
- ▶ The higher are the fourth family members masses, the better are conditions

$$V_{u_1 d_4} > V_{u_1 d_3} ,$$

$$V_{u_2 d_4} < V_{u_1 d_4} , \text{ and}$$

$$V_{u_3 d_4} < V_{u_1 d_4}$$

fulfilled.

- ▶ The **stable family** of the **upper four families** group is the candidate to form the **Dark Matter**.
- ▶ Masses of the upper four families are influenced :
 - by the $\widetilde{SU}(2)_{\parallel \widetilde{SO}(3,1)} \times \widetilde{SU}(2)_{\parallel \widetilde{SO}(4)}$ **scalar fields** with the corresponding family quantum numbers,
 - by the **scalars** $(A_{78}^Q, A_{78}^{Q'}, A_{78}^{Y'})$, and
 - by the **condensate** of the two ν_R of the **upper four families**.


Matter-antimatter asymmetry

There are also **triplet** and **anti-triplet** scalars, $s = (9, \dots, d)$:

	state	τ^{33}	τ^{38}	spin	τ^4	Q
$A_{9\ 10}^{Ai}$ (+)	$A_9^{Ai} - iA_{10}^{Ai}$	$+\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	0	$-\frac{1}{3}$	$-\frac{1}{3}$
$A_{11\ 12}^{Ai}$ (+)	$A_{11}^{Ai} - iA_{12}^{Ai}$	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	0	$-\frac{1}{3}$	$-\frac{1}{3}$
$A_{13\ 14}^{Ai}$ (+)	$A_{13}^{Ai} - iA_{14}^{Ai}$	0	$-\frac{1}{\sqrt{3}}$	0	$-\frac{1}{3}$	$-\frac{1}{3}$
$A_{9\ 10}^{Ai}$ (-)	$A_9^{Ai} + iA_{10}^{Ai}$	$-\frac{1}{2}$	$-\frac{1}{2\sqrt{3}}$	0	$+\frac{1}{3}$	$+\frac{1}{3}$
$A_{11\ 12}^{Ai}$ (-)	$A_{11}^{Ai} + iA_{12}^{Ai}$	$\frac{1}{2}$	$-\frac{1}{2\sqrt{3}}$	0	$+\frac{1}{3}$	$+\frac{1}{3}$
$A_{13\ 14}^{Ai}$ (-)	$A_{13}^{Ai} + iA_{14}^{Ai}$	0	$\frac{1}{\sqrt{3}}$	0	$+\frac{1}{3}$	$+\frac{1}{3}$

They cause transitions from anti-leptons into quarks and anti-quarks into quarks and back, **transforming matter into antimatter and back**. The condensate breaks CP symmetry, offering the explanation for the matter-antimatter asymmetry in the universe.

Let us look at scalar triplets, causing the birth of a proton from the left handed positron, antiquark and quark:



These two quarks, d_R^{c1} and u_R^{c3} can bind (at low enough energy) together with u_R^{c2} into the colour **chargeless baryon - a proton**.

After the appearance of the **condensate** the **CP is broken**.

In the expanding universe, fulfilling the Sakharov request for appropriate non-thermal equilibrium, **these triplet scalars have a chance to explain the matter-antimatter asymmetry**.

The opposite transition makes the proton decay.

These processes seem to explain the lepton number non conservation.

Dark matter

$d \rightarrow (d - 4) + (3 + 1)$ before (or at least at) the electroweak break.

- ▶ We follow the **evolution of the universe**, in particular the **abundance of the fifth family members** - the **candidates** for the **dark matter** in the universe.
- ▶ We estimate the behaviour of our stable heavy family quarks and anti-quarks in the expanding universe by solving the system of **Boltzmann equations**.
- ▶ We follow the **clustering** of the **fifth family** quarks and antiquarks into the **fifth family baryons** through the **colour** phase transition.
- ▶ The **mass** of the fifth family members is determined from the today **dark matter density**.

Phys. Rev. D (2009) 80.083534




Figure: The dependence of the two number densities n_{q_5} (of the fifth family quarks) and n_{c_5} (of the fifth family clusters) as the function of $\frac{m_{q_5} c^2}{T k_b}$ is presented for the values $m_{q_5} c^2 = 71 \text{ TeV}$, $\eta_{c_5} = \frac{1}{50}$ and $\eta_{(q\bar{q})_b} = 1$. We take $g^* = 91.5$.

We estimated from following the fifth family members in the expanding universe:



$$10 \text{ TeV} < m_{q_5} c^2 < 4 \cdot 10^2 \text{ TeV} .$$



$$10^{-8} \text{ fm}^2 < \sigma_{c_5} < 10^{-6} \text{ fm}^2 .$$

(It is at least $10^{-6} \times$ smaller than the cross section for the first family neutrons.)

We estimate from the scattering of the fifth family members on the ordinary matter on our Earth, on the direct measurements - DAMA, CDMS,... - ...



$$200 \text{ TeV} < m_{q_5} c^2 < 10^5 \text{ TeV} .$$

- ▶ In the *standard model* the **family members** with all their properties, the **families**, the **gauge vector fields**, the **scalar Higgs**, the **Yukawa couplings**, exist by the **assumption**.
- ▶ ** In the **spin-charge-family theory** the appearance and all the properties of all these fields follow from the simple starting action with **two kinds of spins** and with the **gravity only** .
 - ** The theory offers the explanation for the **dark matter**.
 - ** The theory offers the explanation for the **matter-antimatter asymmetry**.
 - ** All the **scalar** and all the **vector** gauge fields are **directly or indirectly observable**.
- ▶ ** The **spin-charge-family theory** even offers the **creation and annihilation operators without postulation**.

The *spin-charge-family theory* explains also many other properties, which are not explainable in the *standard model*, like "miraculous" non-anomalous triangle Feynman diagrams.

The more work is put into the *spin-charge-family theory* the more explanations for the phenomena follow.

Concrete predictions:

- ▶ There are several scalar fields;
 - **two triplets** , ◦ **three singlets** ,explaining **higgs** and **Yukawa couplings**, some of them will be observed at the LHC, JMP 6 (2015) 2244, Phys. Rev. D 91 (2015) 6, 065004.
- ▶ There is the **fourth family**, (weakly) coupled to the observed **three**, which will be observed at the LHC, New J. of Phys. 10 (2008) 093002.
- ▶ There is the **dark matter** with the predicted properties, Phys. Rev. D (2009) 80.083534.
- ▶ There is the ordinary **matter/antimatter asymmetry** explained and the **proton decay** predicted and explained, Phys. Rev. D 91 (2015) 6, 065004.

We recognize that:

- ▶ The last **data for mixing matrix of quarks** are in better agreement with our prediction for the **3×3 submatrix** elements of the **4×4 mixing matrix** than the previous ones.
- ▶ Our **fit** to the last data predicts how will the **3×3 submatrix elements change** in the next more accurate measurements.
- ▶ Masses of the **fourth family** lie **much above** the known three, masses of quarks are close to each other.
- ▶ The **larger are masses of the fourth family the larger are $V_{u_1 d_4}$ in comparison with $V_{u_1 d_3}$ and the more is valid that $V_{u_2 d_4} < V_{u_1 d_4}$, $V_{u_3 d_4} < V_{u_1 d_4}$.**
The flavour changing neutral currents are correspondingly weaker.

- ▶ Masses of the **fifth family** lie **much above** the known three and the **predicted fourth family** masses.
- ▶ Although the upper four families carry the weak (of two kinds) and the colour charge, these group of four families are completely decoupled from the lower four families up to the $< 10^{16}$ GeV, unless the breaks of symmetries recover.
- ▶ **Baryons** of the **fifth family** are heavy, forming small enough clusters with small enough scattering amplitude among themselves and with the ordinary matter to be the candidate for the **dark matter**.
- ▶ The "nuclear" force among them is different from the force among ordinary nucleons.

- ▶ The **spin-charge-family** theory is offering an explanation for the **hierarchy problem**:
The mass matrices of the **two four families groups** are almost democratic, causing spreading of the **fermion masses** from 10^{16} GeV to 10^{-8} MeV.

To summarize:

- ▶ I hope that I managed to convince you that I can answer many open questions of particle physics and cosmology.
- ▶ **The collaborators are very welcome!**
- ▶ There are namely a lot of properties to derive.